

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12603 A1

(51) International Patent Classification⁷: C07D 209/80,
495/04, A61K 31/403, 31/407, A61P 25/28

(21) International Application Number: PCT/GB00/03011

(22) International Filing Date: 4 August 2000 (04.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9918962.3 11 August 1999 (11.08.1999) GB

(71) Applicant (for all designated States except US): VER-
NALIS RESEARCH LIMITED [GB/GB]; Oakdene
Court, 613 Reading Road, Winnersh, Wokingham RG41
5UA (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): BENTLEY,
Jonathan, Mark [GB/GB]; Oakdene Court, 613 Reading
Road, Winnersh, Wokingham RG41 5UA (GB). ROFFEY,
Jonathan, Richard, Anthony [GB/GB]; Oakdene Court,
613 Reading Road, Winnersh, Wokingham RG41 5UA
(GB). DAVIDSON, James, Edward, Paul [GB/GB];
Oakdene Court, 613 Reading Road, Winnersh, Woking-
ham RG41 5UA (GB). MANSELL, Howard, Langham
[GB/GB]; Oakdene Court, 613 Reading Road, Winnersh,
Wokingham RG41 5UA (GB). HAMLYN, Richard, John
[GB/GB]; Oakdene Court, 613 Reading Road, Winnersh,
Wokingham RG41 5UA (GB). CLIFFE, Ian, Anthony

[GB/GB]; Oakdene Court, 613 Reading Road, Winnersh,
Wokingham RG41 5UA (GB). ADAMS, David, Reginald
[GB/GB]; Oakdene Court, 613 Reading Road, Winnersh,
Wokingham RG41 5UA (GB). MONCK, Nathaniel,
Julius [GB/GB]; Oakdene Court, 613 Reading Road,
Winnersh, Wokingham RG41 5UA (GB).

(74) Agents: HOWARD, Paul, Nicholas et al.; Carpmals &
Ransford, 43 Bloomsbury Square, London WC1A 2RA
(GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG):

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: INDOLE DERIVATIVES, PROCESS FOR THEIR PREPARATION, PHARMACEUTICAL COMPOSITIONS CON-
TAINING THEM AND THEIR MEDICINAL APPLICATION



(57) Abstract: A chemical compound of formula (I) wherein: R₁ and R₂ are independently selected from hydrogen and alkyl; R₃ is alkyl; R₄, R₆ and R₇ are independently selected from hydrogen, halogen, hydroxy, alkyl, aryl, amino, alkylamino, dialkylamino, alkoxy, aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl, nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl; R₅ is selected from hydrogen, halogen, hydroxy, alkyl, aryl, amino, alkylamino, dialkylamino, alkoxy, aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl, nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl; A is a 5- or 6-membered partially unsaturated or aromatic heterocyclic ring or a 5- or 6-membered partially unsaturated carbocyclic ring, wherein if A is a 6-membered partially unsaturated carbocyclic ring

then at least one of R₄ to R₇ is other than hydrogen, and pharmaceutically acceptable salts, addition compounds and prodrugs thereof, and the use thereof in therapy, particularly as an agonist or antagonist of a 5HT_{2C} receptor, particularly a 5HT_{2C} receptor, for instance in the treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus, and sleep apnea, and particularly for the treatment of obesity.

INDOLE DERIVATIVES, PROCESS FOR THEIR PREPARATION, PHARMACEUTICAL COMPOSITIONS CONTAINING THEM AND THEIR MEDICINAL APPLICATION

The present invention relates to indole derivatives, to processes and intermediates for their preparation, to pharmaceutical compositions containing them and to their medicinal use. The active compounds of the present invention are useful in treating obesity and other disorders.

It has been recognised that obesity is a disease process influenced by environmental factors in which the traditional weight loss methods of dieting and exercise need to be supplemented by therapeutic products (S. Parker, "*Obesity: Trends and Treatments*", *Scrip Reports*, PJB Publications Ltd, 1996).

Whether someone is classified as overweight or obese is generally determined on the basis of their body mass index (BMI) which is calculated by dividing body weight (kg) by height squared (m^2). Thus, the units of BMI are kg/m^2 and it is possible to calculate the BMI range associated with minimum mortality in each decade of life. Overweight is defined as a BMI in the range 25-30 kg/m^2 , and obesity as a BMI greater than 30 kg/m^2 . There are problems with this definition in that it does not take into account the proportion of body mass that is muscle in relation to fat (adipose tissue). To account for this, obesity can also be defined on the basis of body fat content: greater than 25% and 30% in males and females, respectively.

As the BMI increases there is an increased risk of death from a variety of causes that is independent of other risk factors. The most common diseases with obesity are cardiovascular disease (particularly hypertension), diabetes (obesity aggravates the development of diabetes), gall bladder disease (particularly cancer) and diseases of reproduction. Research has shown that even a modest reduction in body weight can correspond to a significant reduction in the risk of developing coronary heart disease.

Compounds marketed as anti-obesity agents include Orlistat (Reductil®) and Sibutramine. Orlistat (a lipase inhibitor) inhibits fat absorption directly and tends to produce a high incidence of unpleasant (though relatively harmless) side-effects such as diarrhoea. Sibutramine (a mixed 5-HT/noradrenaline reuptake inhibitor) can increase blood

pressure and heart rate in some patients. The serotonin releaser/reuptake inhibitors fenfluramine (Pondimin[®]) and dexfenfluramine (Redux[™]) have been reported to decrease food intake and body weight over a prolonged period (greater than 6 months). However, both products were withdrawn after reports of preliminary evidence of heart valve abnormalities associated with their use. There is therefore a need for the development of a safer anti-obesity agent.


The non-selective 5-HT_{2C} receptor agonists/partial agonists m-chlorophenylpiperazine (mCPP) and trifluoromethylphenylpiperazine (TFMPP) have been shown to reduce food intake in rats (G.A. Kennett and G. Curzon, *Psychopharmacol.*, 1988, 98, 93-100; G.A. Kennett, C.T. Dourish and G. Curzon, *Eur. J. Pharmacol.*, 1987, 141, 429-453) and to accelerate the appearance of the behavioural satiety sequence (S.J. Kitchener and C.T. Dourish, *Psychopharmacol.*, 1994, 113, 369-377). Recent findings from studies with mCPP in normal human volunteers and obese subjects have also shown decreases in food intake. Thus, a single injection of mCPP decreased food intake in female volunteers (A.E.S. Walsh *et al.*, *Psychopharmacol.*, 1994, 116, 120-122) and decreased the appetite and body weight of obese male and female subjects during subchronic treatment for a 14 day period (P.A. Sargeant *et al.*, *Psychopharmacol.*, 1997, 113, 309-312). The anorectic action of mCPP is absent in 5-HT_{2C} receptor knockout mutant mice (L.H. Tecott *et al.*, *Nature*, 1995, 374, 542-546) and is antagonised by the 5-HT_{2C} receptor antagonist SB-242084 in rats (G.A. Kennett *et al.*, *Neuropharmacol.*, 1997, 36, 609-620). It seems therefore that mCPP decreases food intake via an agonist action at the 5-HT_{2C} receptor.

Other compounds which have been proposed as 5-HT_{2C} receptor agonists for use in the treatment of obesity include the substituted 1-aminoethyl indoles disclosed in EP-A-0655440. CA-2132887 and CA-2153937 disclose that tricyclic 1-aminoethylpyrrole derivatives and tricyclic 1-aminoethyl pyrazole derivatives bind to 5-HT_{2C} receptors and may be used in the treatment of obesity. WO-A-98/30548 discloses aminoalkylindazole compounds as 5-HT_{2C} agonists for the treatment of CNS diseases and appetite regulation disorders. Substituted 1,2,3,4-Tetrahydrocarbazoles have been reported as synthetic trypanocides in *J. Med. Chem.*, 1970, 13, 327 and *J. Med. Chem.*, 1973, 16, 1411. 9-(2-Dialkylaminopropyl)-1,2,3,4-tetrahydrocarbazoles have been disclosed in US 2687414 and US 2541211. 7-Substituted-9-(2-dialkylaminoethyl)-1,2,3,4-tetrahydrocarbazoles have

been disclosed in DE 930988. The pharmacological behaviour of 2,3-polymethyleneindoles has been described in *J. Med. Chem.*, 1964, 69, 2910. Derivatives of polynuclear indoles have been described as antidepressants in *J. Med. Chem.*, 1964, 7, 625. Amino-substituted penthienoindoles with pharmacological properties are disclosed in US 3142678. 1,2,3,4-Tetrahydro-cyclopent[b]indoles are disclosed in FR 2242983 and DE 2438413. 4-(3-Aminobutyl)-1,2,3,4-tetrahydrocyclopent[b]indole has been described in *Khim. Geterotsikl. Soedin*, 1970, 6, 371.

It is an object of this invention to provide selective, directly acting 5HT₂ receptor
10 ligands for use in therapy and particularly for use as anti-obesity agents. It is a further
object of this invention to provide directly acting ligands selective for 5-HT_{2B} and/or 5-
HT_{2C} receptors, for use in therapy and particularly for use as anti-obesity agents. It is a
further object of this invention to provide selective, directly acting 5-HT_{2C} receptor
ligands, preferably 5-HT_{2C} receptor agonists, for use in therapy and particularly for use as
15 anti-obesity agents.

According to the present invention there is provided a chemical compound of formula (I):



(I)

wherein:

R₁ and R₂ are independently selected from hydrogen and alkyl;

R₃ is alkyl;

R₄, R₆ and R₇ are independently selected from hydrogen, halogen, hydroxy, alkyl, aryl,
25 amino, alkylamino, dialkylamino, alkoxy, aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl,
nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl;

R₅ is selected from hydrogen, halogen, hydroxy, alkyl, aryl, amino, alkylamino, dialkylamino, alkoxy, aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl, nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl; and

A is a 5- or 6- membered partially unsaturated or aromatic heterocyclic ring or a 5- or 6-
5 membered partially unsaturated carbocyclic ring,

wherein if A is a 6-membered partially unsaturated carbocyclic ring then at least one of R₄ to R₇ is other than hydrogen,

and pharmaceutically acceptable salts, addition compounds and prodrugs thereof.

10 As used herein, the term "alkyl" means a branched or unbranched, cyclic or acyclic, saturated or unsaturated (e.g. alkenyl or alkynyl) hydrocarbyl radical. Where cyclic, the alkyl group is preferably C₃ to C₁₂, more preferably C₅ to C₁₀. Where acyclic, the alkyl group is preferably C₁ to C₁₀, more preferably C₁ to C₆, more preferably methyl, ethyl, propyl (n-propyl or isopropyl), butyl (n-butyl, isobutyl or tertiary-butyl) or pentyl (including n-pentyl
15 and iso-pentyl), more preferably methyl. It will be appreciated therefore that the term "alkyl" as used herein includes alkyl (branched or unbranched), alkenyl (branched or unbranched), alkynyl (branched or unbranched), cycloalkyl, cycloalkenyl and cycloalkynyl.

As used herein, the term "lower alkyl" means a branched or unbranched, cyclic or
20 acyclic, saturated or unsaturated (e.g. alkenyl or alkynyl) hydrocarbyl radical, wherein a cyclic lower alkyl group is C₅, C₆ or C₇, and wherein an acyclic lower alkyl group is methyl, ethyl, propyl (n-propyl or isopropyl) or butyl (n-butyl, isobutyl or tertiary-butyl), more preferably methyl.

25 As used herein, the term "aryl" means an aromatic group, such as phenyl or naphthyl, or a heteroaromatic group containing one or more heteroatom, such as pyridyl, pyrrolyl, quinoliny, furanyl, thienyl, oxadiazolyl, thiadiazolyl, thiazolyl, oxazolyl, isoxazolyl, pyrazolyl, triazolyl, imidazolyl or pyrimidinyl.

30 As used herein, the term "alkoxy" means alkyl-O-. As used herein, the term "lower alkoxy" means loweralkyl-O-. As used herein, the term "aryloxy" means aryl-O-.

As used herein, the term "halogen" means a fluorine, chlorine, bromine or iodine radical, preferably a fluorine or chlorine radical.

As used herein the term "prodrug" means any pharmaceutically acceptable prodrug of
5 the compound of formula (I) which is metabolised *in vivo* to a compound of formula (I).

As used herein, the term "pharmaceutically acceptable salt" means any pharmaceutically acceptable salt of the compound of formula (I). Salts may be prepared from pharmaceutically acceptable non-toxic acids and bases including inorganic and organic acids and bases. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric,
10 dichloroacetic, ethanesulfonic, formic, fumaric, gluconic, glutamic, hippuric, hydrobromic, hydrochloric, isethionic, lactic, maleic, malic, mandelic, methanesulfonic, mucic, nitric, oxalic, pantoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, oxalic, p-toluenesulfonic and the like. Particularly preferred are fumaric, hydrochloric, hydrobromic, phosphoric, succinic, sulfuric and methanesulfonic acids, particularly fumaric acid. Acceptable base salts
15 include alkali metal (e.g. sodium, potassium), alkaline earth metal (e.g. calcium, magnesium) and aluminium salts.

As used herein, the term "addition compound" means any pharmaceutically acceptable addition compound of the compound of formula (I). Addition compounds include
20 those which are formed without change of valency from the union between a compound of formula (I) and one or more other molecules, particularly solvates, hydrates and inclusion complexes (such as cyclodextrin complexes).

As used herein, the term "A is a 5- or 6-membered ring" refers to a ring containing 5
25 or 6 ring atoms in total, i.e. including the carbon atoms in the unsaturated positions of the indole ring to which A is fused.

As used herein, the term "carbocyclic ring" refers to a ring wherein all the ring atoms are carbon atoms.

30

As used herein, the term "partially unsaturated ring" refers to a ring which contains unsaturated ring atoms and one or more double bonds but which is not aromatic, for example a cyclohexenyl, cyclopentenyl, or thiacyclohexenyl ring. It will be appreciated therefore that

a partially unsaturated ring A may contain one double bond, i.e. the double bond between the unsaturated 2 and 3 positions of the indole ring to which the ring A is fused, in which case the atoms of the ring A, other than the carbon atoms in the unsaturated 2 and 3 positions of the indole ring to which A is fused, are saturated. Alternatively, a partially unsaturated ring A may contain an additional double bond provided that this additional double bond does not result in the ring A being aromatic.

Where any of R₁ to R₇ is an alkyl group or an alkyl-containing group (such as alkoxy, alkylamino or alkylthio, for instance) as defined in formula (I) above, then that alkyl group, or the alkyl group of the alkyl-containing group, may be substituted or unsubstituted. Where any of R₄ to R₇ is an aryl group or an aryl-containing group (such as aryloxy, for instance) as defined in formula (I), then said aryl group, or the aryl group of the aryl-containing group, may be substituted or unsubstituted. The ring A may be substituted or unsubstituted, preferably unsubstituted. Where any of R₁ to R₇ or A is substituted, there will generally be 1 to 3 substituents present, preferably 1 substituent. Substituents may include:

carbon-containing groups such as

alkyl,

aryl, (e.g. substituted and unsubstituted phenyl),

arylalkyl; (e.g. substituted and unsubstituted benzyl);

halogen atoms and halogen containing groups such as

haloalkyl (e.g. trifluoromethyl),

haloaryl (e.g. chlorophenyl);

oxygen containing groups such as

oxo,

alcohols (e.g. hydroxy, hydroxyalkyl, hydroxyaryl, (aryl)(hydroxy)alkyl),

ethers (e.g. alkoxy, aryloxy, alkoxyalkyl, aryloxyalkyl, alkoxyaryl, aryloxyaryl),

aldehydes (e.g. carboxaldehyde),

ketones (e.g. alkylcarbonyl, arylcarbonyl, alkylcarbonylalkyl, alkylcarbonylaryl, arylcarbonylalkyl, arylcarbonylaryl, arylalkylcarbonyl, arylalkylcarbonylalkyl, arylalkylcarbonylaryl)

- acids (e.g. carboxy, carboxyalkyl, carboxyaryl),
acid derivatives such as esters
(e.g. alkoxycarbonyl, aryloxy carbonyl,
alkoxycarbonylalkyl, aryloxy carbonylalkyl,
5 alkoxycarbonylaryl, aryloxy carbonylaryl,
alkylcarbonyloxy, alkylcarbonyloxyalkyl),
amides
(e.g. aminocarbonyl, mono- or di-alkylaminocarbonyl,
aminocarbonylalkyl, mono- or di-
10 alkylaminocarbonylalkyl, arylaminocarbonyl or
arylalkylaminocarbonyl, alkylcarbonylamino,
arylcarbonylamino or arylalkylcarbonylamino),
carbamates
(eg. alkoxycarbonylamino, aryloxy carbonylamino,
15 arylalkyloxy carbonylamino, aminocarbonyloxy, mono-
or di-alkylaminocarbonyloxy, arylaminocarbonyloxy
or arylalkylaminocarbonyloxy)
and ureas
(eg. mono- or di-alkylaminocarbonylamino,
20 arylaminocarbonylamino or
arylalkylaminocarbonylamino);
nitrogen containing groups such as
amines (e.g. amino, mono- or dialkylamino, arylamino,
aminoalkyl, mono- or dialkylaminoalkyl),
25 azides,
nitriles (e.g. cyano, cyanoalkyl),
nitro;
sulfur containing groups such as
thiols, thioethers, sulfoxides, and sulfones
30 (e.g. alkylthio, alkylsulfinyl, alkylsulfonyl,
alkylthioalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl,
arylthio, arylsulfinyl, arylsulfonyl, arylthioalkyl,
arylsulfinylalkyl, arylsulfonylalkyl)

and heterocyclic groups containing one or more, preferably one, heteroatom,

(e.g. thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, oxadiazolyl, thiadiazolyl, aziridiny, azetidiny, pyrrolidiny, 5 pyrroliny, imidazolidiny, imidazolinyl, pyrazolidiny, tetrahydrofuranyl, pyranyl, pyronyl, pyridyl, pyrazinyl, pyridazinyl, piperidyl, hexahydroazepiny, piperazinyl, morpholinyl, thianaphthyl, benzofuranyl, isobenzofuranyl, indolyl, 10 oxyindolyl, isoindolyl, indazolyl, indolinyl, 7-azaindolyl, benzopyranyl, coumarinyl, isocoumarinyl, quinolinyl, isoquinolinyl, naphthridiny, cinnolinyl, quinazolinyl, pyridopyridyl, benzoxazinyl, quinoxalinyl, 15 chromenyl, chromanyl, isochromanyl, phthalazinyl and carbolinyl).

It is preferred that the compounds of formula (I) are selected from those wherein R_1 to R_7 and A are as defined above with the proviso that if A is a 5- or 6- membered partially 20 unsaturated carbocyclic ring then at least one of R_4 to R_7 is other than hydrogen.

In the compounds of formula (I), preferably R_1 and R_2 are independently selected from hydrogen and lower alkyl (preferably acyclic lower alkyl and more preferably methyl), and preferably from hydrogen. 25

In one embodiment, the compounds of formula (I) are selected from compounds in which R_1 is the same as R_2 . Preferably, R_1 and R_2 are both hydrogen.

The compounds of formula (I) are preferably selected from compounds in which R_3 is 30 lower alkyl, preferably acyclic lower alkyl, and more preferably methyl.

R_5 is selected from hydrogen, halogen, hydroxy, alkyl (including cycloalkyl, halo-alkyl (such as trifluoromethyl) and arylalkyl), aryl, amino, alkylamino, dialkylamino,

alkoxy (including arylalkoxy), aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl, nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl.

5 In one embodiment, R_5 is selected from halogen, hydroxy, alkyl (including cycloalkyl, halo-alkyl (such as trifluoromethyl) and arylalkyl), aryl, amino, alkylamino, dialkylamino, alkoxy (including arylalkoxy), aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl, nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl.

10 Preferably R_5 is selected from hydrogen, halogen and alkoxy, preferably from alkoxy and halogen, and preferably from alkoxy. Where R_5 is halogen, it is preferred that R_5 is selected from fluoro, chloro and bromo, preferably from fluoro and chloro and more preferably from fluoro. Where R_5 is selected from alkoxy, it is preferred that R_5 is selected from lower alkoxy, preferably acyclic lower alkoxy.

15 R_4 , R_6 and R_7 are independently selected from hydrogen, halogen, hydroxy, alkyl (including cycloalkyl, halo-alkyl (such as trifluoromethyl) and arylalkyl), aryl, amino, alkylamino, dialkylamino, alkoxy (including arylalkoxy), aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl, nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl.

20 Preferably R_4 is selected from hydrogen, halogen, alkyl and alkoxy, and is preferably hydrogen. Where R_4 is alkyl, it is preferred that R_4 is lower alkyl, preferably acyclic lower alkyl. Where R_4 is alkoxy, it is preferred that R_4 is lower alkoxy, preferably acyclic lower alkoxy.

25 Preferably R_6 is selected from hydrogen and halogen. Where R_6 is selected from halogen, R_6 is preferably fluoro or chloro, preferably fluoro.

30 Preferably R_7 is selected from hydrogen, halogen and alkoxy, preferably from hydrogen and halogen, and preferably from halogen. Where R_7 is alkoxy, it is preferred that R_7 is lower alkoxy, preferably acyclic lower alkoxy. Where R_7 is halogen, it is preferred that R_7 is selected from fluoro, chloro and bromo, preferably from chloro and bromo and preferably chloro.

It is preferred that at least one of R₄ to R₇ is a group other than hydrogen.

Where A is a heterocyclic ring, A may contain one or more heteroatom(s), and preferably only one heteroatom. Where A contains one or more heteroatom(s), it is preferred
5 that the heteroatoms are selected from N, O and S. Where A is partially unsaturated, it is preferred that A contains no heteroatoms.

It is preferred that A is a 5-membered ring.

10 It is preferred that A is partially unsaturated, preferably wherein the atoms of the ring A, other than the carbon atoms in the unsaturated 2 and 3 positions of the indole ring to which the ring A is fused, are saturated.

In one embodiment, the compounds of formula (I) are selected from compounds
15 wherein A is a 5-membered partially unsaturated carbocyclic ring, a 5-membered heterocyclic ring (preferably aromatic) or a 6-membered partially unsaturated carbocyclic ring, preferably from compounds wherein A is a 5-membered partially unsaturated carbocyclic ring or a 5-membered heterocyclic ring, and more preferably from compounds wherein A is a 5-membered partially unsaturated carbocyclic ring.

20

In a further embodiment, the compounds of formula (I) are selected from compounds wherein A is selected from the group consisting of cyclopentenyl (including oxocyclopentenyl (particularly 1-oxocyclopent-4-enyl)), cyclohexenyl, thiacyclohexenyl (particularly 4-thiacyclohexenyl) and thienyl.

25

The compounds of the invention may contain one or more asymmetric carbon atoms, so that the compounds can exist in different stereoisomeric forms. The compounds can be, for example, racemates or optically active forms. The optically active forms can be obtained by resolution of the racemates or by asymmetric synthesis. In a preferred
30 embodiment of the invention, where all of R₄ to R₇ are hydrogen, the preferred stereochemistry at the carbon atom to which R₃ and NR₁R₂ are bound is (*R*). In an alternative embodiment, where R₅ or R₇ is a group other than hydrogen, the preferred stereochemistry at the carbon atom to which R₃ and NR₁R₂ are bound is (*S*).

In one embodiment of the invention, the compounds of formula (I) are preferably selected from:

- (S)-1-(7,8-difluoro-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine,
- 5 (S)-1-(7-fluoro-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine,
- (S)-1-(8-chloro-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine,
- (S)-1-(6-methoxy-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine,
- (S)-1-(7-fluoro-6-methoxy-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine,
- (S)-1-(7-fluoro-8-methoxy-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine,
- 10 (S)-1-(8-chloro-7-fluoro-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine,
- (S)-1-(1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine and
- (R)-1-(1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine.

According to a further aspect of the invention, there is provided a compound of
15 formula (I) for use in therapy.

The compounds of formula (I) may be used in the treatment (including prophylactic treatment) of disorders associated with 5-HT₂ receptor function. The compounds may act as receptor agonists or antagonists, preferably receptor agonists. Preferably, the
20 compounds may be used in the treatment (including prophylactic treatment) of disorders associated with 5-HT_{2B} and 5-HT_{2C} receptor function. Preferably, the compounds may be used in the treatment (including prophylactic treatment) of disorders where 5-HT_{2C} receptor activity is required, and preferably where a 5HT_{2C} receptor agonist is required.

25 The compounds of formula (I) may be used in the treatment or prevention of central nervous disorders such as depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, personality disorders, age-
30 related behavioural disorders, behavioural disorders associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa or premenstrual tension; damage of the central nervous system such as by trauma,

stroke, neurodegenerative diseases or toxic or infective CNS diseases such as encephalitis or meningitis; cardiovascular disorders such as thrombosis; gastrointestinal disorders such as dysfunction of gastrointestinal motility; diabetes insipidus; and sleep apnea.

5 According to a further aspect of the invention, there is provided use of a compound of formula (I) in the manufacture of a medicament for the treatment (including prophylaxis) of the above-mentioned disorders. In a preferred embodiment, there is provided use of a compound of formula (I) in the manufacture of a medicament for the treatment (including prophylaxis) of obesity.

10

 According to a further aspect of the invention, there is provided a method of treating a disorder selected from the group consisting of the above-mentioned disorders comprising administering to a patient in need of such treatment an effective dose of a compound of formula (I). In a preferred embodiment, there is provided a method of
15 treatment (including prophylaxis) of obesity.

 According to a further aspect of the invention, there is provided a pharmaceutical composition comprising a compound of formula (I) in combination with a pharmaceutically acceptable carrier or excipient and a method of making such a
20 composition comprising combining a compound of formula (I) with a pharmaceutically acceptable carrier or excipient.


 According to a further aspect of the invention, there is provided a method of preparing a compound of formula (I), for instance in the manner described below in the
25 Reaction Schemes. R_1 to R_7 are as previously defined.

 As used herein, the term "saturated 2,3-ring-fused indoles" refers to a tricyclic compound having a ring A as defined herein which is fused to an indole ring across the double bond in the 2- and 3-positions of the indole ring, wherein the atoms of the ring A,
30 other than the carbon atoms in the unsaturated 2- and 3-positions of the indole ring to which A is fused, are saturated.

As used herein, the term "unsaturated 2,3-ring-fused indoles" refers to a tricyclic compound having a ring A as defined herein which is fused to an indole ring across the double bond in the 2- and 3-positions of the indole ring, wherein one or more of the atoms of the ring A, other than the carbon atoms in the unsaturated 2- and 3-positions of the indole ring to which A is fused, are unsaturated. It will be understood that the term "unsaturated 2,3-ring-fused indoles" includes compounds wherein the ring A is aromatic.

In Reaction Scheme 1, the saturated 2,3-ring-fused indoles (IV) may be formed by sequential reaction of the suitably substituted N-2-bromophenyl acetamide (eg R = CF₃) (II) with methyllithium and the appropriate 2-halo-cyclic ketone (III), followed by *tert* butyllithium and then trifluoroacetic acid. The N-alkyl ring-fused indole (V) (eg R = *tert* Bu) may then be obtained by reaction of (IV) with an appropriate carbamylethylsulfonate in the presence of a strong base such as potassium hydroxide in a solvent such as methyl sulfoxide. The indole (I) (R₁ = R₂ = H) may then be obtained by reaction of the indole (V) with a reagent suitable to reveal the protected amine function.

Reaction Scheme 1



The compounds of formula (I) (R_1 and/or R_2 = alkyl) may be prepared from compounds of formula (I) ($R_1 = R_2 = H$) by standard methods such as reductive alkylation with an appropriate aldehyde or ketone in the presence of a reducing agent such as sodium triacetoxyborohydride, formic acid or sodium cyanoborohydride.


5

The unsaturated 2,3-ring-fused indoles (I) may be formed in a similar manner to the saturated 2,3-ring-fused indoles (I), through the intermediacy of the unsaturated 2,3-ring-fused indole (IV) obtained from the saturated 2,3-ring-fused indole (IV) under standard dehydrogenation conditions such as through treatment with DDQ or Pd on carbon in a
10 suitable solvent such as dioxan and xylene respectively.

Alternatively, compounds of the invention can be conveniently prepared according to Reaction Scheme 2. Treatment of phenylhydrazine (II) with a cyclic ketone under acidic conditions in a suitable solvent, such as ethanol or water, produces indole (III).

15 Reaction of indole (III) with an alkylating agent such as *tert*-butyl [2-[(1-methanesulfonyl)oxy]propyl]carbamate in the presence of a base such as potassium hydroxide in a suitable solvent *e.g.* methyl sulfoxide gives indole-carbamate (IV). A compound of formula (I) where $R_1 = R_2 = H$ can be prepared by treatment of (IV) with an acid such as hydrochloric acid in a suitable solvent such as methanol or by use of a strong
20 base such as potassium *tert*-butoxide in a solvent such as methyl sulfoxide. A compound of formula (I) where R_1 and / or R_2 = alkyl can be prepared by reductive alkylation using an aldehyde or ketone in the presence of a reducing agent such as formic acid, sodium cyanoborohydride or sodium triacetoxyborohydride.

Reaction Scheme 2



- If, in any of the other processes mentioned herein, the substituent groups R_1 , R_2 , R_3 , R_4 , R_5 , R_6 or R_7 is other than the one required, the substituent group may be converted to the desired substituent by known methods. The substituents R_1 , R_2 , R_3 , R_4 , R_5 , R_6 or R_7 may also need protecting against the conditions under which the reaction is carried out. In such a case, the protecting group may be removed after the reaction has been completed.
- The processes described above may be carried out to give a compound of the invention in the form of a free base or as an acid addition salt. If the compound of the invention is obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid addition salt. Conversely, if the product of the process is a free base, an acid addition salt may be obtained by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from basic compounds.

The compositions of the present invention may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers. Thus, the active compounds of the invention may be formulated for oral, buccal, intranasal, parenteral (e.g.,

intravenous, intramuscular or subcutaneous) transdermal or rectal administration or in a form suitable for administration by inhalation or insufflation.

For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (*e.g.* pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropylmethylcellulose); fillers (*e.g.* lactose, microcrystalline cellulose or calcium phosphate); lubricants (*e.g.* magnesium stearate, talc or silica); disintegrants (*e.g.* potato starch or sodium starch glycollate); or wetting agents (*e.g.* sodium lauryl sulfate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (*e.g.* sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agents (*e.g.* lecithin or acacia); non-aqueous vehicles (*e.g.* almond oil, oily esters or ethyl alcohol); and preservatives (*e.g.* methyl or propyl *p*-hydroxybenzoates or sorbic acid).

For buccal administration the composition may take the form of tablets or lozenges formulated in conventional manner.

The active compounds of the invention may be formulated for parenteral administration by injection, including using conventional catheterization techniques or infusion. Formulations for injection may be presented in unit dosage form *e.g.* in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulating agents such as suspending, stabilising and/or dispersing agents.

Alternatively, the active ingredient may be in powder form for reconstitution with a suitable vehicle, *e.g.* sterile pyrogen-free water, before use.

The active compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other glycerides.

5 For intranasal administration or administration by inhalation, the active compounds of the invention are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped by the patient or as an aerosol spray presentation from a pressurized container or a nebulizer, with the use of a suitable propellant, *e.g.* dichlorodifluoromethane, trichlorofluoromethane,
10 dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the
15 invention and a suitable powder base such as lactose or starch.

A proposed dose of the active compounds of the invention for oral, parenteral or buccal administration to the average adult human for the treatment of the conditions referred to above (*e.g.*, obesity) is 0.1 to 500 mg of the active ingredient per unit dose
20 which could be administered, for example, 1 to 4 times per day.

The invention will now be described in detail with reference to the following examples. It will be appreciated that the invention is described by way of example only and modification of detail may be made without departing from the scope of the invention.
25

EXPERIMENTAL

Assay Procedures

30 1. Binding to serotonin receptors

The binding of compounds of formula (I) to serotonin receptors was determined *in vitro* by standard methods. The preparations were investigated in accordance with the assays given hereinafter.

Method (a): For the binding to the 5-HT_{2c} receptor the 5-HT_{2c} receptors were radiolabelled with [³H]-5-HT. The affinity of the compounds for 5-HT_{2c} receptors in a CHO cell line was determined according to the procedure of D. Hoyer, G. Engel and H.O. Kalkman, *European J. Pharmacol.*, 1985, 118, 13-23.

Method (b): For the binding to the 5-HT_{2B} receptor the 5-HT_{2B} receptors were radiolabelled with [³H]-5-HT. The affinity of the compounds for human 5-HT_{2B} receptors in a CHO cell line was determined according to the procedure of K. Schmuck, C. Ullmer, P. Engels and H. Lubbert, *FEBS Lett.*, 1994, 342, 85-90.

Method (c): For the binding to the 5-HT_{2A} receptor the 5-HT_{2A} receptors were radiolabelled with [¹²⁵I]-DOI. The affinity of the compounds for 5-HT_{2A} receptors in a CHO cell line was determined according to the procedure of D. J. McKenna and S. J. Peroutka, *J. Neurosci.*, 1989, 9/10, 3482-90.

The thus determined activity of compounds of formula (I) is shown in Table 1.

Table 1: Radioligand Binding Data

Compound	K _i (2C) / nM	K _i (2A) / nM	K _i (2B) / nM
Example 1	65	122	40
Example 11	63	314	210
Example 14	64	375	180
Example 26	106	144	127
Example 27	141	545	496
Example 29	474	823	653
Example 30	19	48	31
Example 31	65	550	161
Example 32	27	106	58
Example 33	63	233	152
Example 37	41	86	65
Example 43	62	167	162

2. Functional activity

The functional activity of compounds of formula (I) was assayed using a
5 Fluorimetric Imaging Plate reader (FLIPR) in the following manner.

CHO cells expressing either the h5-HT_{2C} or h5-HT_{2A} receptors were counted and
plated into standard 96 well microtitre plates before the day of testing to give a confluent
monolayer. The following day the cells were dye loaded with the calcium sensitive dye
Fluo 3-AM by incubation with serum free culture maintenance media containing pluronic
10 acid and Fluo 3-AM dissolved in DMSO at 37 °C in a CO₂ incubator at 95% humidity for
approximately 90 minutes. Unincorporated dye was removed by washing with Hanks
balanced salt solution containing 20mM HEPES and 2.5mM probenecid (the assay buffer)
using an automated cell washer to leave a total volume of 100 µL/well.

The drug (dissolved in 50 µL of assay buffer) was added at a rate of 70 µL/sec to
15 each well of the FLIPR 96 well plate during fluorescence measurements. The
measurements are taken at 1 sec intervals and the maximum fluorescent signal was
measured (approx 10-15 secs after drug addition) and compared with the response
produced by 10 µM 5-HT (defined as 100%) to which it is expressed as a percentage
response (relative efficacy). Dose response curves were constructed using Graphpad Prism
20 (Graph Software Inc.).

The thus determined activity of compounds of formula (I) is shown in Table 2.

Table 2: Functional Data


Compound	h5-HT _{2A}		h5-HT _{2C}	
	EC ₅₀ (nM)	Relative Efficacy (%)	EC ₅₀ (nM)	Relative Efficacy (%)
Example 1	10000	0	272	77
Example 2	10000	0	347	85
Example 4	10000	60	179	65
Example 11	1686	25	89	85
Example 14	6247	48	252	80
Example 15	10000	0	1732	93
Example 16	10000	0	307	86
Example 18	2102	63	36	75
Example 30	361	43	90	72
Example 33	10000	22	316	81
Example 36	1339	25	189	64
Example 37	2990	28	127	84
Example 42	805	51	87	74

5

Synthetic Examples

Example 1: (S)-1-(7,8-Difluoro-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine fumarate

10



2'-Bromo-2,2,2-trifluoroacetanilide

To a stirred solution of 2-bromo-4,5-difluoroaniline [H. Ishikawa, T. Uno, H. Miyamoto, H. Hiraki, H. Tamaoka, M. Tominaga and K. Nakagawa, *Chem. Pharm. Bull.*, 1990, 38(9), 2459-2462] (7.2 g, 34 mmol) in ether (50 mL) at 0 °C was added sodium carbonate (5.4 g, 44 mmol) and trifluoroacetic anhydride (6.2 mL, 44 mmol). The reaction mixture was stirred at room temperature for 1 h. Water (100 mL) was added and the mixture was extracted with dichloromethane (3 x 100 mL). The organic extracts were combined, dried (magnesium sulfate), filtered and concentrated *in vacuo* to give the product (9.9 g, 94%) as a white solid. IR ν_{\max} (Nujol)/cm⁻¹ 3270, 1716, 1550, 1489, 1465, 1226, 1181, 919, 876 and 821; NMR δ_{H} (400 MHz, CDCl₃) 7.45-7.5 (1H, dd, *J* 7.5 Hz), 8.28-8.34 (1H, dd, *J* 8 Hz) and 8.36 (1H, br s).

7,8-Difluoro-1,2,3,3a,4,8a-hexahydro-8a-hydroxy-cyclopent[b]indole

A stirred solution of 2'-Bromo-2,2,2-trifluoroacetanilide (5.3 g, 35 mmol), in tetrahydrofuran (200 mL) was cooled to -78 °C. A solution of methyllithium (12.5 mL, 35 mmol, 1.4 M in ether) was added maintaining the temperature of reaction below -75 °C. After 10 min a solution of *tert*-butyllithium (20.5 mL, 70 mmol, 1.7 M in pentane) was added over 5 min and the reaction was stirred for 1 h at -78 °C. The mixture was warmed to -50 °C and 2-chlorocyclopentanone (2.1 mL, 42 mmol) was added dropwise. The reaction was warmed slowly to room temperature and stirred for a further 2 h. A solution of potassium hydroxide in methanol (10%, 20 mL) was added and the mixture was stirred at room temperature for 12 h. The mixture was poured onto dilute hydrochloric acid (5%, 150 mL) and washed with dichloromethane (3 x 150 mL). The aqueous layer was basified (15% aqueous sodium hydroxide solution) and extracted with dichloromethane (3 x 150 mL). The organic extracts were combined, dried (magnesium sulfate), filtered and concentrated *in vacuo* to give the product (0.85 g, 11%) as a pale brown solid. *R*_f 0.39 [SiO₂; heptane-ethyl acetate (10:3)]; NMR δ_{H} (400 MHz, CDCl₃) 1.53-1.67 (2H, m), 1.78-1.89 (1H, m), 2.02-2.17 (2H, m), 2.29-2.37 (1H, m), 4.04 (1H, dd, *J* 6 Hz), 6.21-6.26 (1H, m) and 6.86-6.94 (1H, m).

7,8-Difluoro-1,2,3,4-tetrahydrocyclopent[b]indole

A stirred solution of 7,8-difluoro-1,2,3,3a,4,8a-hexahydro-8a-hydroxy-cyclopent[b]indole (1.1 g, 5.2 mmol), in dichloromethane (150 mL) was cooled to 0 °C. Trifluoroacetic acid (20 drops) was added and the reaction mixture was stirred at room temperature for 18 h.

5 The reaction mixture was poured onto saturated sodium hydrogen carbonate solution (20 mL) and extracted with dichloromethane (3 x 50 mL). The organic extracts were combined, dried (magnesium sulfate), filtered, concentrated *in vacuo* and purified by column chromatography [SiO₂; ethyl acetate-heptane (1:5)] to give the product (0.78 g, 78%) as a white crystalline solid. IR ν_{\max} (Nujol)/cm⁻¹ 3467, 2925, 2854, 1565, 1515, 1450,

10 1348, 1327, 1244, 1053, 1025, 977, 857, 783, 630 and 516; NMR δ_{H} (400 MHz, CDCl₃) 2.49-2.58 (2H, m), 2.79-2.87 (2H, m), 2.9-2.96 (2H, m), 6.81-6.95 (2H, m), and 7.83 (1H, br s).

(*S*)-4-[2-(*tert*-Butoxycarbonylamino)propyl]-7,8-difluoro-1,2,3,4-
15 tetrahydrocyclopent[b]indole

7,8-Difluoro-1,2,3,4-tetrahydrocyclopent[b]indole (0.56 g, 2.9 mmol) was added portionwise to a mixture of methyl sulfoxide (15 mL) and crushed potassium hydroxide (0.57 g, 10.2 mmol). The mixture was warmed to 35 °C and stirred for 30 min. A solution

20 of (*S*)-2-(*tert*-butoxycarbonylamino)propane methanesulfonate (1.85 g, 7.3 mmol) in methyl sulfoxide (5 mL) was added over a 1 h period, the mixture was then stirred at 35 °C for 20 h. Water (30 mL) was added and the mixture was extracted with ether (3 x 50 mL). The organic extracts were combined, dried (magnesium sulfate), filtered, concentrated *in vacuo* and purified by column chromatography [SiO₂; heptane-ethyl acetate (5:1)] to give

25 the product (0.55 g, 52%) as a white crystalline solid; IR ν_{\max} (Nujol)/cm⁻¹ 3366, 1684, 1516, 1456, 1248, 1022 and 773; NMR δ_{H} (400 MHz, CDCl₃) 1.1 (3H, d, *J* 7 Hz), 1.43 (9H, br s), 2.48-2.57 (2H, m), 2.79-2.87 (2H, m), 2.91-2.98 (2H, m), 3.84-3.92 (1H, dd, *J* 7 Hz), 3.96-4.07 (1H, m), 4.08 (1H, br s), 4.4 (1H, br s), 6.83-6.92 (1H, m) and 6.94-7.08 (1H, br s).

30

(*S*)-1-(7,8-Difluoro-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine fumarate

A solution of (S)-4-[2-(*tert*-butoxycarbonylamino)propyl]-7,8-difluoro-1,2,3,4-tetrahydrocyclopent[*b*]indole (0.4 g, 1.1 mmol) and trifluoroacetic acid (5 mL) in dichloromethane (15 mL) was stirred at room temperature for 1 h. The mixture was made basic by the addition of aqueous sodium hydroxide solution (2 N), then extracted with
5 dichloromethane (3 x 50 mL). The organic extracts were combined, dried (magnesium sulfate), filtered and concentrated *in vacuo* to give an orange oil. The oil was dissolved in 2-propanol (5 mL) and the solution was heated to boiling then fumaric acid (0.38 g, 3.3 mmol) was added. The mixture was cooled to room temperature and filtered. The filter-cake was washed (2-propanol, ether) and dried *in vacuo* to give the title compound (0.89 g,
10 68%) as a pale orange solid. mp. 154-156 °C (dec.); NMR δ_H (400 MHz, DMSO-*d*₆) 1.13 (3H, d, *J* 7 Hz), 2.43-2.52 (2H, m), 2.78-2.94 (4H, m), 3.5-3.57 (1H, m), 4.13 (1H, d, *J* 8 Hz), 4.29 (1H, dd, *J* 6.5 Hz), 6.55 (2H, s), 7.01-7.10 (1H, m) and 7.26-7.31 (1H, m).

Reference herein to (S)-1-(7,8-Difluoro-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine fumarate will be understood to mean a compound prepared by the above
15 synthetic procedure.

Other compounds of formula (I) as defined herein may be prepared according to the following synthetic methods.

20

Phenylhydrazine preparation (General Method A)


Commercially available substituted phenylhydrazines were used with the exception of the compounds listed below in Table 3. The compounds listed in Table 3 were synthesised in accordance with the method (general synthetic method A) given below for compounds 36a,
25 37a and 42a.

Compounds 36a, 37a and 42a: 4-Fluoro-3-methoxyphenylhydrazine hydrochloride

To stirred hydrochloric acid (100 mL) at 0 °C was added 3-methoxy-4-fluoroaniline (10 g,
30 71 mmol) followed by water (10 mL) and more hydrochloric acid (10 mL). The mixture was warmed to room temperature, stirred for 20 min then cooled to -5 °C. A solution of sodium nitrite (5.14 g, 75 mmol) in water (25 mL) was added dropwise such that the internal temperature remained below 0 °C. The mixture was warmed to room temperature

and stirred for 2 h. The mixture was cooled to -5°C and a solution of tin(II)chloride dihydrate (64 g, 284 mmol) in hydrochloric acid (200 mL) was added dropwise such that the internal temperature remained below 0°C . The mixture was warmed to room temperature, stirred for 3 h then filtered. The filter-cake was washed with hydrochloric acid and dried under vacuum to give a pink solid (7.4 g). The precipitate from the combined filtrates was filtered-off, washed (hydrochloric acid) and dried under vacuum to give a further crop of product (1.8 g. to give a combined yield of 9.2 g, 67%). Data for 4-fluoro-3-methoxyphenylhydrazine hydrochloride are included in Table 3 below.

10 Table 3: Phenylhydrazines (prepared by General Method A)



In this structural formula there may be a plurality of R groups, as detailed in Table 3 below.

Compound	R	Yield	Data
23a	3-OBn	72%	Hydrochloride. NMR (400 MHz, CDCl_3) δ_{H} 7.43 (2H, d, J 7.5 Hz), 7.38 (2H, t, J 7.5 Hz), 7.32 (1H, t, J 7 Hz), 7.13 (1H, t, J 8 Hz), 6.48 (1H, t, J 2.5 Hz), 6.45 (1H, dd, J 8, 2.5 Hz), 6.41 (1H, dd, J 8, 2.5 Hz), 5.04 (2H, s); HPLC [Supelcosil ABZ+, 1.0 ml/min, methanol-10 mM aqueous ammonium acetate solution (80:20)] 90% (2.62 min).

24a	3-O ⁱ Pr	52%	Hydrochloride. NMR (400 MHz, DMSO- <i>d</i> ₆) δ_{H} 10.24 (3H, br s, NH ₃), 8.26 (1H, br s, NH), 7.16 (1H, t, <i>J</i> 8.2 Hz), 6.61 (1H, t, <i>J</i> 2.1 Hz), 6.54 (1H, dd, <i>J</i> 8.0, 1.6 Hz), 6.50 (1H, dd, <i>J</i> 8.3, 2.0 Hz), 4.57 (1H, quint, <i>J</i> 6.0 Hz), 1.27 (6H, d, <i>J</i> 6.0 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 90% (2.55 min).
25a	3-O ⁱ Pr	as 24a	as Compound 24a
28a	2-OCF ₃	77%	Hydrochloride. NMR (400 MHz, DMSO- <i>d</i> ₆) δ_{H} 10.56 (3H, br s, NH ₃), 8.41 (1H, br s, NH), 7.37-7.31 (3H, m), 7.03 (1H, dq, <i>J</i> 8.6, 4.3 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99% (2.38 min).
29a	4-OCF ₃	84%	m.p. 216 °C; Found: C, 34.04; H, 3.42; N, 11.11%. C ₇ H ₇ F ₃ N ₂ O.1.5H ₂ O requires: C, 34.06; H, 3.47; N, 11.35%.
33a	3,4-di-F	70%	NMR (400MHz, CDCl ₃) δ_{H} 3.56 (2H, br s), 5.13 (1H, br s), 6.47 (1H, m), 6.69 (1H, m), 6.99 (1H, dd, <i>J</i> 8.53Hz, 17.57Hz); IR ν_{max} (nujol)/cm ⁻¹ 3258, 1613, 1516, 1465, 1265, 1222 and 771.
36a	4-F, 3-OMe	67%	m.p. 250+ °C (dec.); NMR: (400 MHz, DMSO- <i>d</i> ₆) δ_{H} 10.17 (3H, s, NH ₃), 8.14 (1H, s, NH), 7.15 (1H, dd, <i>J</i> 11.6, 8.6 Hz), 6.95 (1H, dd, <i>J</i> 7.6, 3.0 Hz), 6.54 (1H, dt, <i>J</i> 8.6, 3.0 Hz), 3.83 (3H, s, MeO).
37a	4-F, 3-Ome	as 36a	as Compound 36a
42a	4-F, 3-Ome	as 36a	as Compound 36a

Fischer Synthesis of indoles (General Method B)

The indoles listed in Table 4 below were synthesised in accordance with the following synthetic methods (General Methods B(i) and B(ii)) given below for compounds 14b, 30b, 11b and 12b

5 **Method B(i): Aqueous Sulfuric Acid**

Compounds 14b and 30b: 1,2,3,4-Tetrahydrocyclopent[*b*]indole

A solution of phenylhydrazine (32.44 g, 300 mmol) in 2-propanol (300 mL) was treated
10 with cyclopentanone (27 mL, 25.7 g, 305 mmol). The solution was stirred at 20 °C for 1 h
and poured onto a mixture of ice (900 g) and water (300 mL). The chilled mixture was
stirred until the ice melted and then filtered. The filter-cake was washed with water (2 x
300 mL) to give an off-white, moist solid (85 g). The solid was added to water (540 mL)
and the stirred suspension was treated with concentrated sulfuric acid (33 mL, 61 g, 600
15 mmol). The suspension was then heated under reflux for 30 min, cooled to 0 °C and then
stirred for 15 min. The dark-red solid was filtered off, washed with water (2 x 60 mL) and
air-dried for 18 h. The crude product was added to stirred dichloromethane (300 mL),
stirred for 30 min and filtered. The tarry residue was washed with dichloromethane (100
mL) and the filtrate was treated with silica (48 g), stirred for 1 h and filtered. The silica
20 residue was washed with dichloromethane (400 mL) and the filtrate was concentrated to
give a solid, which was triturated with hexane to give 1,2,3,4-tetrahydrocyclopent[*b*]indole
(30 g, 65%) as a pink solid. Analytical data for 1,2,3,4-tetrahydrocyclopent[*b*]indole are
included in Table 4 below.

25 Where the intermediate hydrazone was obtained as an oil the following method was used:

A solution of the arylhydrazine (100 mmol) in benzene (100 mL) was treated with
cyclopentanone (9 mL, 8.6 g, 102 mmol). The solution was heated under reflux with
azeotropic removal of water for 30-60 min. The solution was allowed to cool and was
30 concentrated *in vacuo* to give the arylhydrazone as an oil which was used directly in the
subsequent step as described above.

Method B(ii): Ethanol as solvent


Compounds 11b and 12b: 1,2,3,4-Tetrahydro-6-methoxy-cyclopent[*b*]indole and 1,2,3,4-tetrahydro-8-methoxy-cyclopent[*b*]indole


- 5 To stirred, degassed ethanol (20 mL), shielded from light and under an atmosphere of Ar at ambient temperature, was added 3-methoxyphenylhydrazine hydrochloride (1.0 g, 5.6 mmol) and cyclopentanone (0.5 mL, 5.7 mmol). The mixture was heated at reflux for 24 h, cooled to room temperature then poured onto 300 mL ice-water and made basic with saturated aqueous sodium bicarbonate solution (to pH 8). The suspension was filtered, and
- 10 the resultant solid was washed with water and dried to afford the crude product as a dark brown solid (0.95 g, 89%) which was purified by flash column chromatography [SiO_2 ; isohexane-dichloromethane (3:2 \rightarrow 1:1)] afforded the separated isomeric indole products. Alternatively the crude product was purified by filtration of a dichloromethane solution through a plug of silica and concentration *in vacuo* followed by trituration with toluene,
- 15 filtration, and washing of the resultant solid with ice-cold toluene-heptane (1:1) to afford exclusively the 6-isomer. Analytical data for 1,2,3,4-tetrahydro-6-methoxy-cyclopent[*b*]indole and 1,2,3,4-tetrahydro-8-methoxy-cyclopent[*b*]indole are included in Table 4 below.
- 20 For the appropriate examples, pentindole regioisomers arising from the use of unsymmetrical arylhydrazines were separated by column chromatography, recrystallisation from toluene, cyclohexane, isohexane or ethanol or by trituration with toluene or pentane.

Table 4: Indoles formed using General Methods B(i) and B(ii)


25

In this structural formula, there may be an additional double bond in the 5- or 6-membered ring fused to the indole ring. In Table 4 below, the substituents R_4 to R_7 are hydrogen unless otherwise stated in column 2.




Compound	Substitution Pattern (method)	n	Yield	Data
2b	R ₆ =F (i)	1	67%	m.p. 102-103 °C (Ethanol); Found: C, 75.36; H, 5.80; N, 7.97%. C ₁₁ H ₁₀ FN requires: C, 75.41; H, 5.75; N, 7.99%.
3b	R ₅ =Cl (i)	2	18%	m.p. 181 °C (Ethanol); Found: C, 70.03; H, 5.87; N, 6.85%. C ₁₂ H ₁₂ ClN requires: C, 70.07; H, 5.88; N, 6.81%.
4b	R ₇ =Cl (i)	1	23%	Low-melting solid from mother liquors of 6-chloro isomer recrystallisation. NMR (400 MHz, CDCl ₃) δ _H 7.88 (1H, m, NH), 7.16 (1H, dd, <i>J</i> 1, 8 Hz), 7.03 (1H, dd, <i>J</i> 1, 8 Hz), 6.96 (1H, t, <i>J</i> 8 Hz), 3.04 (2H, tt, <i>J</i> 1.5, 7 Hz), 2.85 (2H, tt, <i>J</i> 1.5, 7 Hz), 2.53 (2H, quint., <i>J</i> 7 Hz); HPLC: [Supelcosil ABZ+; 1.0 mL/min, methanol-10 mM aqueous ammonium acetate solution (80:20)] 80% (8.00 min) + 6-chloro isomer (20%).
5b	R ₅ =Cl (i)	1	21%	m.p. 188-191 °C (Ethanol); Found: C, 69.21; H, 5.18; N, 7.31% C ₁₁ H ₁₀ ClN requires: C, 68.94; H, 5.26; N, 7.30%.
6b	R ₅ = Cl; synthetic method is (ii); n in the above formula is not applicable; the compound contains an S-heteroatom: 		37%	m.p. 179-182 °C (Ethanol); Found: C, 59.29; H, 4.44; N, 6.28; S, 14.38; Cl, 16.04%. C ₁₁ H ₁₀ CINS requires: C, 59.06; H, 4.51; N, 6.26; S, 14.33; Cl, 15.85%.

7b	R ₅ =Br (i)	1	12%	m.p. 199.5-200 °C (dec.); Found: C, 55.48; H, 4.21; N, 5.85%. C ₁₁ H ₁₀ BrN.0.125H ₂ O requires: C, 55.43; H, 4.33; N, 5.86%.
8b	R ₅ =Br (i)	2	3.4%	NMR (400 MHz, CDCl ₃) δ _H 7.67 (1H, m, NH), 7.41 (1H, d, <i>J</i> 1.5 Hz), 7.30 (1H, d, <i>J</i> 8.5 Hz), 7.16 (1H, dd, <i>J</i> 1.5, 8.5 Hz), 2.73-2.64 (4H, m), 1.95-1.82 (4H, m); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99% (10.12 min).
9b	R ₆ =Cl (i)	2	35%	NMR (400 MHz, CDCl ₃) δ _H 7.67 (1H, m, NH), 7.40 (1H, d, <i>J</i> 2 Hz), 7.16 (1H, d, <i>J</i> 8.5 Hz), 7.04 (1H, dd, <i>J</i> 2, 8.5 Hz), 2.74-2.69 (2H, m), 2.67-2.63 (2H, m), 1.94-1.82 (4H, m); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99% (9.28 min).
10b	R ₆ =Cl (i)	1	42%	NMR (400 MHz, CDCl ₃) δ _H 7.84 (1H, m, NH), 7.39 (1H, d, <i>J</i> 2 Hz), 7.19 (1H, d, <i>J</i> 8.5 Hz), 7.03 (1H, dd, <i>J</i> 8.5, 2 Hz), 2.86 (2H, m), 2.79 (2H, tt, <i>J</i> 6.5, 1.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99% (7.67 min).
11b	R ₅ =OMe (ii)	1	30%	m.p. 136-137.5 °C; NMR (400 MHz, CDCl ₃) δ _H 7.68 (1H, m, NH), 7.29 (1H, d, <i>J</i> 8.5 Hz), 6.81 (1H, d, <i>J</i> 2 Hz), 6.74 (1H, dd, <i>J</i> 2, 8.5 Hz), 3.83 (3H, s), 2.85-2.76 (4H, m), 2.55-2.47 (2H, m).

12b	$R_7=OMe$ (ii)	1		m.p. 87-89 °C; NMR (400 MHz, $CDCl_3$) δ_H 7.79 (1H, m, NH), 6.99 (1H, t, J 8 Hz), 6.91 (1H, dd, J 8, 1 Hz), 6.49 (1H, d, J 8 Hz), 3.90 (3H, s), 2.98-2.93 (2H, m), 2.84-2.78 (2H, m), 2.55-2.47 (2H, m).
13b	$R_4=R_5=Cl$ (i)	1	28%	m.p. 104-107 °C (isohexane); Found: C, 58.65; H, 4.04; N, 6.20; Cl, 31.30%. $C_{11}H_9Cl_2N$ requires: C, 58.43; H, 4.01; N, 6.19; Cl, 31.36%.
14b	(i)	1	65%	m.p. 107-108 °C (hexane); Found: C, 83.04; H, 7.12; N, 8.78%. $C_{11}H_{11}N \cdot 0.1H_2O$ requires: C, 83.09; H, 7.10; N, 8.81%.
15b	$R_5 = R_7 = Cl$; synthetic method is (ii); n in the above formula is not applicable; the compound contains an S-heteroatom: 		7%	(Synthesised using tetrahydrothiophen-3-one, initial product aromatises during reaction) m.p. 105 °C (heptane); NMR (400 MHz, $CDCl_3$) δ_H 8.20 (1H, m, NH), 7.44 (1H, d, J 5.5 Hz), 7.26 (1H, d, J 1.5 Hz), 7.17 (1H, d, J 1.5 Hz), 7.03 (1H, d, J 5.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (90:10)] 99% (6.66 min).
16b	$R_5=Cl$; $R_6=F$ (i)	1	21%	m.p. 139.5-140 °C (cyclohexane); Found: C, 62.87; H, 4.35; N, 6.69%. $C_{11}H_9ClFN$ requires: C, 63.02; H, 4.33; N, 6.68%.
17b	$R_5=CF_3$ (i)	1	33%	m.p. 161-162 °C (pentane); Found: C, 63.87; H, 4.46; N, 6.18%. $C_{12}H_{10}FN$ requires: C, 64.00; H, 4.48; N, 6.22%.

18b	R ₇ =Cl; R ₆ =F (i)	1	40%	Low-melting solid. NMR (400 MHz, CDCl ₃) δ_H 7.86 (1H, m, NH), 7.07 (1H, dd, <i>J</i> 3.5, 9 Hz), 6.86 (1H, t, <i>J</i> 9 Hz), 3.03 (2H, tt, <i>J</i> 1.5, 7 Hz), 2.84 (2H, t, <i>J</i> 7 Hz) and 2.53 (2H, quintet, <i>J</i> 7 Hz); HPLC: [Xterra; 2.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99.5% (8.29 min).
19b	R ₅ =R ₆ =Cl (i)	1	12%	m.p. 169 °C (Toluene); Found: C, 58.45; H, 3.95; N, 6.19%. C ₁₁ H ₉ Cl ₂ N requires: C, 58.43; H, 4.01; N, 6.19%.
20b	R ₆ =OMe (i*)	1	85%	(*as method (i) but at room temperature and in water). NMR (400 MHz, CDCl ₃) δ_H 7.71 (1H, m, NH), 7.18 (1H, d, <i>J</i> 8.5 Hz), 6.91 (1H, d, <i>J</i> 2.5 Hz), 6.74 (1H, dd, <i>J</i> 8.5, 2.5 Hz), 3.85 (3H, s), 2.87-2.78 (4H, m), 2.56-2.49 (2H, m); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 94% (3.81 min).
21b	R ₇ =CF ₃ (i)	1	50%	NMR (400 MHz, CDCl ₃) δ_H 8.05 (1H, m, NH), 7.44 (1H, d, <i>J</i> 8 Hz), 7.37 (1H, d, <i>J</i> 8 Hz), 7.12 (1H, t, <i>J</i> 8 Hz), 2.95-2.87 (4H, m), 2.60-2.50 (2H, m); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99% (6.63 min).
22b	R ₆ =R ₇ =Cl (i)	1	6%	(Mixture with 7,8-dichloro product). m.p. 107-114 °C; HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 50% (12.25 min).

23b	R ₇ =OBn (ii)	1	10%	NMR (400 MHz, CDCl ₃) δ_H 7.79 (1H, m, NH), 7.50 (2H, d, <i>J</i> 7.5 Hz), 7.38 (2H, t, <i>J</i> 7.5 Hz), 6.97 (1H, t, <i>J</i> 8 Hz), 6.93 (1H, dd, <i>J</i> 8, 1 Hz), 6.56 (1H, dd, <i>J</i> 8, 1 Hz), 5.18 (2H, s), 3.01 (2H, t, <i>J</i> 7 Hz), 2.83 (2H, t, <i>J</i> 7 Hz), 2.52 (2H, quint., <i>J</i> 7 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (9.24 min).
24b	R ₇ =O ⁱ Pr (ii)	1	4%	NMR (400 MHz, CDCl ₃) δ_H 7.74 (1H, br s, NH), 6.94 (1H, t, <i>J</i> 7.8 Hz), 6.88 (1H, dd, <i>J</i> 8.2, 0.9 Hz), 6.50 (1H, d, <i>J</i> 6.9 Hz), 4.57 (1H, quint, <i>J</i> 6.0 Hz), 2.93 (2H, obs tt, <i>J</i> 6.9, 1.5 Hz), 2.80 (2H, obs tt, <i>J</i> 6.5, 1.5 Hz), 2.52-2.45 (2H, m), 1.34 (6H, d, <i>J</i> 6.0 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 77% (4.87 min), material decomposes under mildly acidic conditions.
25b	R ₅ =O ⁱ Pr (ii)	1	2%	MS [Found: (m/z) = 215. C ₁₄ H ₁₇ NO requires: M ⁺ 215]; HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 40% (4.62 min), material decomposes under mildly acidic oxygenated conditions.
26b	R ₅ =R ₇ =Cl (i)	1	51%	m.p. 61-62 °C (hexane); Found: C, 58.28; H, 3.99; N, 6.28%. C ₁₁ H ₉ Cl ₂ N requires: C, 58.43; H, 4.01; N, 6.19%.

28b	$R_4=OCF_3$ (i)	1	55%	NMR (400 MHz, $CDCl_3$) δ_H 8.07 (1H, br s, NH), 7.32 (1H, d, J 7.5 Hz), 7.01 (1H, t, J 7.6 Hz), 6.96 (1H, dt, J 7.6, 1.3 Hz), 2.86 (2H, obs dd, J 7.9, 6.3 Hz), 2.80 (2H, obs tt, J 7.9, 1.5 Hz), 2.57-2.50 (2H, m); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99% (6.11 min).
29b	$R_6=OCF_3$ (i)	1	89%	NMR (400 MHz, $CDCl_3$) δ_H 7.89 (1H, m, NH), 7.27 (1H, m), 7.23 (1H, d, J 8.5 Hz), 6.95 (1H, dd, J 9, 2 Hz), 2.86 (2H, t, J 7 Hz), 2.81 (2H, t, J 7 Hz), 2.53 (2H, quint., J 7 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99% (6.87 min).
30b	(i)	1	as 14b	as Compound 14b
31b	$R_5=F$ (i)	1	10%	m.p. 128-131 °C (cyclohexane); Found: C, 75.39; H, 5.80; N, 7.98%. $C_{11}H_{10}FN$ requires: C, 75.41; H, 5.75; N, 7.99%.
32b	Synthetic method is (i); N in the above formula is not applicable; the compound contains an S-heteroatom: 		26%	m.p. 153 °C (dec.); Found: C, 67.97; H, 5.08; N, 7.90%. $C_{10}H_9NS \cdot 0.1H_2O$ requires: C, 67.84; H, 5.24; N, 7.91%.
33b	$R_5=R_6=F$ (ii)	1	52%	Mixture of inseparable regioisomers, used without further purification.

34b	R ₇ =Cl, R ₆ =Me (i)	1	25%	Low-melting solid; NMR (400 MHz, CDCl ₃) δ_H 7.61 (1H, m, NH), 6.97 (1H, d, <i>J</i> 8 Hz), 6.87 (1H, d, <i>J</i> 8 Hz), 3.01 (2H, tt, <i>J</i> 1.5, 7 Hz), 2.75 (2H, tt, <i>J</i> 1.5, 7 Hz), 2.47 (2H, m) and 2.41 (3H, s). HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 90% (8.90 min) [and 6-chloro-7-methyl 10% (8.54 min)].
35b	R ₇ =Cl, R ₆ =Me (i)	1	as 34b	as Compound 34b
36b	R ₆ =F, R ₅ =OMe (ii)	1	44%	NMR (400 MHz, DMSO- <i>d</i> ₆) δ_H 10.69 (1H, s, NH), 7.08 (1H, d, <i>J</i> 12.0 Hz), 6.98 (1H, d, <i>J</i> 7.6 Hz), 3.83 (3H, s, MeO), 2.79 (2H, m), 2.69 (2H, t, <i>J</i> 7.0 Hz), 2.50 (2H, m).
37b	R ₆ =F, R ₅ =OMe (ii)	1	as 36b	as Compound 36b
39b	R ₅ =Cl; R ₆ =F (i)	1	as 16b	as Compound 16b
40b	R ₇ =Cl; R ₆ =F (i)	1	as 18b	as Compound 18b
41b	R ₇ =Br (i)	1	37%	NMR (400 MHz, CDCl ₃) δ_H 7.82 (1H, s, NH), 7.19 (1H, d, <i>J</i> 8 Hz), 7.16 (1H, d, <i>J</i> 8 Hz), 6.89 (1H, t, <i>J</i> 8 Hz), 3.08-3.03 (2H, m), 2.88-2.77 (2H, m), 2.55-2.46 (2H, m); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 95% (2.33 min).

42b	R ₆ =F, R ₇ =OMe (ii)	1	-	Material obtained by column chromatography of the mother liquor from Examples 38b and 39b. The material was used immediately without further purification or analysis.
43b	R ₄ =Cl (i)	1	20%	m.p. 64-66 °C (Ethanol – water); Found: C, 68.81; H, 5.24; N, 7.32%. C ₁₁ H ₁₀ CIN requires: C, 68.94; H, 5.26; N, 7.30%.
44b	R ₄ =Cl (i)	1	as 43b	as Compound 43b

Compound 27b: 6-Ethylthio-1,2,3,4-tetrahydrocyclopent[*b*]indole

5 *1,2,3,4-Tetrahydro-6-(triisopropylsilyl)thio-cyclopent-[b]-indole*

Palladium dibenzylidene-acetone (0.155 g, 5 mol%) and tricyclohexylphosphine (0.19 g, 20 mol%), were weighed out into a flask pre-flushed with argon, and subsequently flushed with argon for 5 min before dissolution in toluene (20 mL). The deep red mixture was stirred at room temperature for 5 minutes under argon, then 6-bromo-1,2,3,4-tetrahydrocyclopent[*b*]indole (0.8 g, 3.4 mmol) was added in one portion. After a further 5 min a solution of potassium (triisopropylsilyl)sulfide (*Tetrahedron Letts.*, 1994, 35(20), 3221-3224 and 3225-6)) in tetrahydrofuran (6 mL) was added via syringe over 4 min. The mixture was stirred for 45 min at room temperature, heated at 80 °C (bath temp) for 70 min then cooled to room temperature over 16 h. The mixture was partitioned between toluene (40 mL) and water (60 mL). The separated aqueous layer was extracted with toluene (30 mL) and the combined organic extracts were washed with brine (20 mL), dried (Na₂SO₄), filtered and concentrated *in vacuo*. The residue was purified by column chromatography [SiO₂; heptane-ethyl acetate (98:2) to (96:4)] to yield 1,2,3,4-tetrahydro-6-(triisopropylsilyl)thio cyclopent[*b*]indole as a pale yellow solid (0.85 g, 73%); NMR (400 MHz, CDCl₃) δ_H 7.76 (1H, br s, NH), 7.43 (1H, d, *J* 1.5 Hz), 7.27-7.25 (1H, m), 7.18 (1H, dd, *J* 8.2, 1.5 Hz), 2.85 (2H, obs dt, *J* 6.9, 1.6 Hz), 2.79 (2H, obs t, *J* 7.0 Hz), 2.52 (2H, obs quint, *J* 7.0 Hz), 1.29-1.19 (3H, m), 1.08 (18H, d, *J* 7.0 Hz); HPLC: [Supelcosil ABZ+;

1.0 ml/min, methanol-10mM aqueous ammonium acetate solution, (90:10)] 99% (11.1min).

6-Ethylthio-1,2,3,4-tetrahydrocyclopent[b]indole

5

A solution of 1,2,3,4-tetrahydro-6-triisopropylsilylthio-cyclopent-[b]-indole (439 mg, 1.31 mmol) and cesium fluoride (395 mg, 2.62 mmol) in dimethyl formamide was stirred at room temperature for 30 min. Iodoethane (0.21 mL, 2.62 mmol) was added dropwise to the suspension and the reaction was stirred at room temperature for 16 h. The reaction
10 mixture was poured onto ice-water (50 mL) and then extracted with ethyl acetate (3 x 50 mL). The organic extracts were combined, dried (magnesium sulfate) and concentrated *in vacuo*. The residue was purified by column chromatography [SiO₂; heptane – ethyl acetate (5:1)] to afford the title compound (158 mg, 56%) as a white solid; NMR (400MHz, CDCl₃) δ_H 1.26 (3H, t, *J* 7.03Hz), 2.53 (2H, m), 2.78-2.93 (6H, m), 7.15 (1H, dd, *J* 1.51Hz,
15 8.03Hz), 7.39 (1H, d, *J* 1.51Hz), 7.81 (1H, br s); IR ν_{max} (nujol)/cm⁻¹ 3403, 3382, 2925, 2854, 1456, 1376 and 808.

Indole Alkylation (General Method C)

The indoles prepared in accordance with the above synthetic methods may be alkylated in
20 accordance with the general synthetic method (General Method C) given below for compound 30c. Table 5 gives details of the compounds prepared in this way


Compound 30c: (*R*) 4-[2-(*tert*-Butoxycarbonylamino)propyl]-1,2,3,4-tetrahydrocyclopent[b]indole

25

Methyl sulfoxide (40 mL) was warmed to 40 °C for 15 min and treated with powdered potassium hydroxide (85%, 2.64 g, 40 mmol). The suspension was stirred for 5 min and then 1,2,3,4-tetrahydrocyclopent[b]indole (1.57 g, 10 mmol) was added. The suspension was stirred at 40 °C for 60 min, then a solution of (*R*)-*tert*-butyl [2-[(1-methanesulfonyl)oxy]propyl]carbamate (6.33 g, 25 mmol) in methyl sulfoxide (13 mL)
30 was added dropwise in portions every 10 min over 90 min. The resultant suspension was stirred at 40 °C for 18 h and then cooled. Di-*tert*-butyl dicarbonate (2.3 mL, 2.2 g, 10 mmol) was added and the suspension was stirred for a further 2 h at 20 °C and poured onto


a mixture of ice (165 g) and water (55 mL). The suspension was stirred for 1 h and then the crude product was filtered-off, washed with water (2 x 25 mL) and air-dried for 5 min [alternatively, the work-up employed ethyl acetate extraction and chromatography (SiO₂; ethyl acetate - dichloromethane (0:1 → 1:19)]. The crude product was dissolved in ethyl acetate, dried (magnesium sulfate) and concentrated to give a solid which was triturated with hexane to give the product as an off-white solid (2.34 g, 74%). Data for (*R*) 4-[2-(*tert*-butoxycarbonylamino)propyl]-1,2,3,4-tetrahydrocyclopent[*b*]indole are listed in Table 5.

10 Table 5: Indole-carbamates synthesised in accordance with General Method C




In this structural formula, there may be an additional double bond in the 5- or 6-membered ring fused to the indole ring. In Table 5 below, the substituents R₄ to R₇ are hydrogen unless otherwise stated (see column 2). In Table 5 below, the stereochemistry at the side chain is indicated in column 3.

Compound	Substitution pattern	n	Yield	Data
2c	R ₆ =F	1 (S)	79%	m.p. 169-170 °C (cyclohexane, 2-propanol); Found: C, 68.61; H, 7.68; N, 8.39%. C ₁₉ H ₂₅ FN ₂ O ₂ requires: C, 68.65; H, 7.58; N, 8.42%.
3c	R ₅ =Cl	2 (S)	83%	m.p. 165-166 °C (ethanol); Found: C, 66.16; H, 7.53; N, 7.72%. C ₂₀ H ₂₇ ClN ₂ O ₂ requires: C, 66.19; H, 7.50; N, 7.72%.
4c	R ₇ =Cl	1 (S)	78%	NMR (400 MHz, CDCl ₃) δ _H 7.22 (1H, m), 7.01 (1H, dd, <i>J</i> 1.5, 8 Hz), 6.98 (1H, t, <i>J</i> 8 Hz), 4.39 (1H, m, NH), 4.16 (1H, m), 4.03 (1H, sept., <i>J</i> 7 Hz), 3.92 (1H, q, <i>J</i> 7 Hz), 3.06 (2H, t, <i>J</i> 7 Hz), 2.85 (2H, t, <i>J</i> 7 Hz), 2.52 (2H, quint., <i>J</i> 7 Hz), 1.42 (9H, s), 1.10 (3H, d, <i>J</i> 7 Hz); HPLC [Xterra, 2.0 mL/min; methanol-10 mM aqueous ammonium acetate solution (50:50) to (80:20) over 4 min then (80:20)] 94% (7.87 min).
5c	R ₅ =Cl	1 (S)	94%	m.p. 172-174 °C; NMR (400 MHz, CDCl ₃) δ _H 7.29 (1H, m) 7.29 (1H, d, <i>J</i> 8 Hz), 7.01 (1H, dd, <i>J</i> 1.5, 8 Hz), 4.42 (1H, m, NH), 4.12-3.89 (3H, m), 2.85 (2H, t., <i>J</i> 7 Hz), 2.81 (2H, t, <i>J</i> 7 Hz), 2.52 (2H, quint., <i>J</i> 7 Hz), 1.42 (9H, s), 1.11 (3H, d, <i>J</i> 6.5 Hz).


6c	$R_5 = \text{Cl}$; n in the above formula is not applicable; the compound contains an S-heteroatom:  Stereochemistry is (S)	-	Not isolated, material deprotected <i>in situ</i> with excess potassium hydroxide.
7c	$R_5 = \text{Br}$	1 (S)	43% NMR (400 MHz, CDCl_3) δ_{H} 7.44 (1H, m), 7.25 (1H, d, J 8 Hz), 7.15 (1H, dd, J 8, 1.5 Hz), 4.42 (1H, m, NH), 4.14-3.90 (3H, m), 2.83 (4H, obs. quint., J 7 Hz), 2.52 (2H, quint., J 7 Hz), 1.43 (9H, s), 1.12 (3H, d, J 7 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99% (8.07 min).
8c	$R_5 = \text{Br}$	2 (S)	24% NMR (400 MHz, CDCl_3) δ_{H} 7.45 (1H, m), 7.29 (1H, d, J 8 Hz), 7.14 (1H, dd, J 8, 1.5 Hz), 4.42 (1H, m, NH), 4.11 (1H, m), 4.02 (1H, obs. sept., J 7 Hz), 3.87 (1H, q, J 7 Hz), 2.68 (4H, q, J 6 Hz), 1.96-1.88 (2H, m), 1.88-1.80 (2H, m), 1.40 (9H, s), 1.10 (3H, d, J 6.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (10.12 min).

9c	R ₆ =Cl	2 (S)	30%	NMR (400 MHz, CDCl ₃) δ _H 7.40 (1H, d, <i>J</i> 2 Hz), 7.29 (1H, m), 7.07 (1H, dd, <i>J</i> 8.5, 2 Hz), 4.42 (1H, m, NH), 4.18 (1H, m), 4.02 (1H, dq, <i>J</i> 20, 7 Hz), 3.85 (dd, <i>J</i> 14.5, 7.5 Hz), 2.72 (2H, obs. t, <i>J</i> 6 Hz), 2.66 (2H, obs. t, <i>J</i> 6 Hz), 1.96-1.89 (2H, m), 1.88-1.81 (2H, m), 1.42 (9H, s), 1.08 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (50:50)] 96% (9.58 min).
10c	R ₆ =Cl	1 (S)	29%	NMR (400 MHz, CDCl ₃) δ _H 7.37 (1H, br. d, <i>J</i> 2 Hz), 7.26 (1H, m), 7.04 (1H, dd, <i>J</i> 8.5, 2 Hz), 4.41 (1H, m, NH), 4.16 (1H, m), 4.03 (1H, m), 3.92 (1H, q, <i>J</i> 7 Hz), 2.86 (2H, t, <i>J</i> 7 Hz), 2.81 (2H, t, <i>J</i> 7 Hz), 2.52 (2H, quint., <i>J</i> 7 Hz), 1.44 (9H, s), 1.09 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (7.82 min).
11c	R ₅ =OMe	1 (S)	64%	NMR (400 MHz, CDCl ₃) δ _H 7.28 (1H, d, <i>J</i> 8.5 Hz), 6.94 (1H, m), 6.73 (1H, dd, <i>J</i> 2.5, 8.5 Hz), 4.48 (1H, m, NH), 4.12 (1H, m), 4.05 (1H, m), 3.88 (1H, dd, <i>J</i> 6.5, 14 Hz), 3.87 (3H, s), 2.86-2.78 (4H, m), 2.55-2.46 (2H, m), 1.43 (9H, s), 1.11 (3H, d, <i>J</i> 7 Hz); HPLC: [Supelcosil ABZ+; 1.0 mL/min, methanol-10 mM aqueous ammonium acetate solution (80:20)] 96% (3.87 min).

12c	R ₇ =OMe	1 (S)	95%	NMR (400 MHz, CDCl ₃) δ_H 7.01 (1H, t, <i>J</i> 7.5 Hz), 6.98 (1H, m), 6.48 (1H, dd, <i>J</i> 7, 1 Hz), 4.44 (1H, m, NH), 4.14 (1H, m), 4.04 (1H, m), 3.91 (1H, m), 3.90 (3H, s), 2.97 (2H, t, <i>J</i> 7 Hz), 2.82 (2H, t, <i>J</i> 7 Hz), 2.49 (2H, quint., <i>J</i> 7 Hz), 1.44 (9H, s), 1.09 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+; 1.0 mL/min, methanol-10 mM aqueous ammonium acetate solution (80:20)] 90% (4.44 min).
13c	R ₄ =R ₅ =Cl	1 (S)	87%	m.p. 205-206 °C (cyclohexane, toluene); Found: C, 59.72; H, 6.34; N, 7.29; Cl, 18.77%. C ₁₉ H ₂₄ Cl ₂ N ₂ O ₂ requires: C, 59.54; H, 6.31; N, 7.30; Cl, 18.50%.
14c		1 (S)	51%	m.p. 172-173 °C (isopropyl ether); Found: C, 71.46; H, 8.22; N, 8.78%. C ₁₉ H ₂₆ N ₂ O ₂ ·0.25H ₂ O requires: C, 71.55; H, 8.38; N, 8.78%.
15c	R ₅ = R ₇ = Cl; n in the above formula is not applicable; the compound contains an S-heteroatom  Stereochemistry is (S)		82%	m.p. 201 °C (hexane); Found: C, 53.53; H, 4.99; N, 6.90%. C ₁₈ H ₂₀ Cl ₂ N ₂ O ₂ ·0.25H ₂ O requires: C, 53.60; H, 5.12; N, 6.95%.
16c	R ₅ =Cl R ₆ =F	1 (S)	74%	m.p. 173.5-176 °C (hexane); Found: C, 61.45; H, 6.54; N, 7.49%. C ₁₉ H ₂₄ ClFN ₂ O ₂ ·0.25H ₂ O requires: C, 61.45; H, 6.65; N, 7.54%.
17c	R ₅ =CF ₃	1 (S)	76%	147-151 °C (hexane); Found: C, 62.22; H, 6.70; N, 7.24%. C ₂₀ H ₂₅ F ₃ N ₂ O ₂ ·0.25H ₂ O requires: C, 62.08; H, 6.64; N, 7.24%.

18c	R ₇ =Cl R ₆ =F	1 (S)	88%	m.p. 161-162 °C (2-propanol); NMR (400 MHz, CDCl ₃) δ _H 7.17 (1H, m), 6.88 (1H, t, <i>J</i> 9 Hz), 4.40 (1H, m), 4.17 (1H, m), 4.01 (1H, dt, <i>J</i> 7, 12.5 Hz), 3.89 (1H, q, <i>J</i> 7 Hz), 3.05 (2H, t, <i>J</i> 7 Hz), 2.84 (2H, t, <i>J</i> 7 Hz), 2.52 (2H, quintet, <i>J</i> 7 Hz), 1.42 (9H, s) and 1.10 (3H, d, <i>J</i> 6.5 Hz).
19c	R ₅ =R ₆ =Cl	1 (S)	81%	m.p. 183-184 °C (hexane); Found: C, 59.45; H, 6.29; N, 7.25%. C ₁₉ H ₂₄ Cl ₂ N ₂ O ₂ requires: C, 59.54; H, 6.31; N, 7.30%.
20c	R ₆ =OMe	1 (S)	63%	m.p. 121 °C; Found: C, 69.66; H, 8.36; N, 7.94%. C ₂₀ H ₂₈ N ₂ O ₃ requires: C, 69.74; H, 8.19; N, 8.13%.
21c	R ₇ =CF ₃	1 (S)	62%	m.p. 154-155 °C (hexane); Found: C, 61.71; H, 6.60; N, 7.13%. C ₂₀ H ₂₅ F ₃ N ₂ O ₂ ·0.5H ₂ O requires: C, 61.37; H, 6.70; N, 7.16%.
22c	R ₆ =R ₇ =Cl	1 (S)	65%	m.p. 152-154 °C (hexane); Found: C, 59.01; H, 6.27; N, 7.08%. C ₁₉ H ₂₄ Cl ₂ N ₂ O ₂ ·0.25H ₂ O requires: C, 58.84; H, 6.37; N, 7.22%.
23c	R ₇ =OBn	1 (S)		NMR (400 MHz, CDCl ₃) δ _H 7.49 (2H, d, <i>J</i> 7 Hz), 7.38 (2H, t, <i>J</i> 7 Hz), 7.30 (1H, t, <i>J</i> 7 Hz), 6.99 (2H, m), 6.55 (1H, m), 5.18 (2H, s), 4.44 (1H, m, NH), 4.15 (1H, m), 4.05 (1H, obs. septet, <i>J</i> 6.5 Hz), 3.92 (1H, q, <i>J</i> 7 Hz), 3.02 (2H, t, <i>J</i> 7 Hz), 2.84 (2H, t, <i>J</i> 7 Hz), 2.51 (2H, quint., <i>J</i> 7 Hz), 1.44 (9H, s), 1.10 (3H, d, <i>J</i> 7 Hz); HPLC: [Supelcosil ABZ+, 1.0 mL/min, methanol-10 mM aqueous ammonium acetate solution (80:20)] 97% (9.80 min).
24c	R ₇ =O ^t Pr	1 (S)	4%	Used immediately without purification or characterisation.
25c	R ₅ =O ^t Pr	1 (S)	2%	Used immediately without purification or characterisation.

26c	$R_5=R_7=Cl$	1 (S)	75%	m.p. 166-166.5 °C (hexane); Found: 58.90; H, 6.22; N, 7.16%. $C_{19}H_{24}Cl_2N_2O_2 \cdot 0.25H_2O$ requires: C, 58.84; H, 6.37; N, 7.22%.
27c	$R_5=EtS$	1 (S)	33%	NMR (400MHz, $CDCl_3$) δ_H 1.11 (3H, d, J 6.02Hz), 1.25 (3H, t, J 6.53Hz), 1.41 (9H, br s), 2.5 (2H, m), 2.77-2.94 (6H, m), 3.91-4.16 (3H, m), 7.12 (1H, d, J 7.53Hz), 7.32 (1H, d, J 7.53), 7.4 (1H, br s); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 91% (7.61 min).
28c	$R_4=OCF_3$	1 (S)	78%	NMR (400 MHz, $CDCl_3$) δ_H 7.28 (1H, dd, J 6.2, 2.9 Hz), 6.96-6.95 (2H, m), 4.39 (1H, br s), 4.15 (2H, br s), 4.00 (1H, br d, J 6.4 Hz), 2.87 (2H, br s), 2.79 (2H, obs t, J 7.2 Hz), 2.50 (2H, obs t, J 6.7 Hz), 1.32 (9H, br s), 1.13 (3H, br d, J 6.4 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 94% (8.79 min).
29c	$R_6=OCF_3$	1 (S)	30%	m.p. 123 °C; NMR (400 MHz, $CDCl_3$) δ_H 7.32 (1H, d, J 8 Hz), 7.25 (1H, d), 6.96 (1H, dd, J 8, 2 Hz), 4.41 (1H, m, NH), 4.18 (1H, m), 4.04 (1H, sept., J 7 Hz), 3.93 (1H, q, J 7 Hz), 2.90-2.80 (4H, m), 2.53 (2H, quint, J 7 Hz), 1.42 (9H, s), 1.12 (3H, d, J 6.5 Hz), HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 96% (7.47 min).
30c		1 (R)	74%	m.p. 170-172 °C (hexane); Found: C, 71.08; H, 8.27; N, 8.71%. $C_{19}H_{26}N_2O_2 \cdot 0.67H_2O$ requires: C, 71.22; H, 8.39; N, 8.74%.
31c	$R_5=F$	1 (S)	59%	m.p. 167-174 °C (hexane); Found: C, 65.76; H, 7.30; N, 7.98%. $C_{19}H_{25}FN_2O_2 \cdot 0.75H_2O$ requires: C, 65.97; H, 7.72; N, 8.10%.

32c	n in the above formula is not applicable; the compound contains an S-heteroatom:  Stereochemistry is (S)		21%	(aromatisation during reaction and work-up) m.p. 200 °C (hexane); Found: C, 65.08; H, 6.65; N, 8.39%. C ₁₈ H ₂₂ N ₂ O ₂ S requires: C, 65.43; H, 6.71; N, 8.47%.
33c	R ₅ =R ₆ =F	1 (S)	43%	Mixture of inseparable regioisomers (with 7,8-difluoro).
34c	R ₇ =Cl R ₆ =Me	1 (S)	70%	NMR (400 MHz, CDCl ₃) δ _H 7.12 (1H, d, <i>J</i> 8 Hz), 6.92 (1H, d, <i>J</i> 8 Hz), 4.40 (1H, m, NH), 4.14 (1H, m), 4.02 (1H, dt, <i>J</i> 6.5, 12 Hz), 3.90 (1H, q, <i>J</i> 7 Hz), 3.06 (2H, t, <i>J</i> 7 Hz), 2.83 (2H, t, <i>J</i> 7 Hz), 2.50 (2H, quintet, <i>J</i> 7 Hz), 2.42 (3H, s), 1.43 (9H, s) and 1.08 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+; 1.0 mL/min, methanol-10 mM aqueous ammonium acetate solution (80:20)] 98% (8.70 min).
35c	R ₇ =Cl R ₆ =Me	1 (R)	92%	NMR (400 MHz, CDCl ₃) δ _H 7.12 (1H, d, <i>J</i> 8 Hz), 6.91 (1H, d, <i>J</i> 8 Hz), 4.41 (1H, m, NH), 4.12 (1H, m), 4.02 (1H, m), 3.98 (1H, q, <i>J</i> 7 Hz), 3.06 (2H, t, <i>J</i> 7 Hz), 2.82 (2H, t, <i>J</i> 7 Hz), 2.50 (2H, quintet, <i>J</i> 7 Hz), 2.42 (3H, s), 1.43 (9H, s) and 1.08 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 98% (8.62 min).

36c	R ₆ =F R ₅ =OMe	1 (R)	40%	Crystallised from Ethanol/water (5:1); NMR (400 MHz, CDCl ₃) δ _H 7.05 (2H, d, <i>J</i> 12.2 Hz), 4.48-4.34 (1H, m), 4.2-3.98 (2H, m), 3.92 (3H, s, MeO), 3.84 (1H, dd, <i>J</i> 14.0, 7.1 Hz), 2.80 (2H, t, <i>J</i> 7.0 Hz), 2.76 (2H, t, <i>J</i> 7.2 Hz), 2.48 (2H, m), 1.40 (9H, br s), 1.09 (3H, d, <i>J</i> 6.5 Hz);). HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (70:30)] 99% (8.82 min) and [Xterra; 2.0 mL/min, methanol-10 mM aqueous ammonium acetate solution, gradient elution 50% to 80% methanol over the first 4 min, then 80:20] 96% (6.89 min).
37c	R ₆ =F R ₅ =OMe	1 (S)	36%	NMR (400 MHz, CDCl ₃) δ _H 7.08 (1H, br. s), 7.07 (1H, d, <i>J</i> 12 Hz), 4.41 (1H, m, NH), 4.16 (1H, m), 4.12 (1H, m), 3.94 (3H, s), 4.04 (1H, dt, <i>J</i> 6.5, 12 Hz), 3.84 (1H, q, <i>J</i> 7 Hz), 2.80 (4H, m), 2.50 (2H, quintet, <i>J</i> 7 Hz), 1.42 (9H, s), 1.11 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Xterra; 2.0 ml/min, gradient elution, methanol-10 mM aqueous ammonium acetate solution (50:50) to (80:20) over 4 min then (80:20)] 97% (6.33 min).
39c	R ₅ =R ₆ =F	1 (R)	58%	m.p. 176-176.5 °C (hexane); Found: C, 61.71; H, 6.59; N, 7.49%. C ₁₉ H ₂₄ ClFN ₂ O ₂ ·0.25H ₂ O requires: C, 61.45; H, 6.65; N, 7.54%.
40c	R ₇ =Cl R ₆ =F	1 (R)	64%	m.p. 160-161 °C (hexane); Found: C, 62.00; H, 6.61; N, 7.56%. C ₁₉ H ₂₄ ClFN ₂ O ₂ requires: C, 62.21; H, 6.59; N, 7.63%.
41c	R ₇ =Br	1 (S)	29%	m.p. 178 °C (2-propanol); Found: C, 58.02; H, 6.45; N, 7.09%. C ₁₉ H ₂₅ BrN ₂ O ₂ requires: C, 58.02; H, 6.41; N, 7.12%.

42c	R ₆ =F R ₇ =OMe	1 (S)	69%	NMR (400 MHz, CDCl ₃) δ_H 6.99-6.94 (1H, m), 6.84 (1H, dd, <i>J</i> 11.3, 9.4 Hz), 4.44-4.37 (1H, m, NH), 4.16-4.00 (2H, m), 4.00 (3H, s), 3.87 (1H, dd, <i>J</i> 14.0, 7.2 Hz), 2.96 (2H, obs t, <i>J</i> 6.6 Hz), 2.83 (2H, obs t, <i>J</i> 7.3 Hz), 2.51 (2H, quintet, <i>J</i> 7.0 Hz), 1.42 (9H, br s), 1.11 (3H, d, <i>J</i> 6.8 Hz); HPLC: [Xterra; 2.0 mL/min, methanol-10 mM aqueous ammonium acetate solution, gradient elution 50% to 80% over the first 4 min, then 80:20] 99.7% (6.55 min).
43c	R ₄ =Cl	1 (R)	31%	m.p. 193-194 °C; Found: C, 65.27; H, 7.24; N, 7.96%. C ₁₉ H ₂₅ ClN ₂ O ₂ requires: C, 65.41; H, 7.22; N, 8.03%.
44c	R ₄ =Cl	1 (S)	25%	m.p. 192-193 °C; NMR (400 MHz, CDCl ₃) δ_H 7.27 (1H, dd, <i>J</i> 1, 8 Hz), 7.04 (1H, dd, <i>J</i> 1, 8 Hz), 6.93 (1H, t, <i>J</i> 8 Hz), 4.80-4.40 (3H, m), 4.20-4.00 (2H, m), 2.89 (2H, m), 2.81 (2H, t, <i>J</i> 7 Hz), 2.51 (2H, quint., <i>J</i> 7 Hz), 1.28 (9H, s), 1.17 (3H, d, <i>J</i> 6.5 Hz).

Compound 45c: (S) 4-[2-(*tert*-Butoxy-carbonylamino)propyl]-1-oxo-1,2,3,4-tetrahydrocyclopent[*b*]indole

5

To a solution of TEMPO.tetrafluoroborate (2.8g, 11.5mmol) in acetonitrile/water (9:1, 50 mL) was added dropwise a solution of (S) 4-[2-(*tert*-butoxy-carbonylamino)propyl]-1,2,3,4-tetrahydrocyclopent[*b*]indole (1.5 g, 5.1 mmol) in acetonitrile - water (9:1, 50 mL). The mixture was stirred for 16 h., then the solvent was removed *in vacuo* and the residue
 10 adsorbed onto alumina (20 g) and purified by column chromatography [Al₂O₃; heptane - ethyl acetate (10:3)] to afford the product (0.7 g, 42%) as a white solid; NMR (400MHz, DMSO-*d*₆) δ_H 1.13 (3H, d, *J* 6.53Hz), 1.21 (9H, br s), 2.82 (2H, m), 3.09 (2H, m), 3.83-4.28 (3H, m), 6.94 (1H, d, *J* 8.03Hz), 7.19 (1H, t, *J* 7.53Hz), 7.27 (1H, d t, *J* 1.0Hz,

7.53Hz), 7.58-7.71 (2H, m); IR ν_{\max} (nujol)/ cm^{-1} 3365, 2924, 2854, 1685, 1538, 1524, 1478, 1452, 1366, 1248, 1168, 1052 and 743.

Deprotection of the Amine (General Method D)

- 5 The protected amines prepared as described above were deprotected in accordance with the following synthetic methods (General Methods D(i), D(ii) and D(iii)) given below for Examples 23, 36 and 45, to give compounds of formula (I). Data for these Examples are given in Table 6.

10 *Method D(i): Deprotection using Hydrogen Chloride*

Example 23: (S)-1-(8-Benzyloxy-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine, hydrochloride

- 15 To a stirred solution of (S) 8-benzyloxy-4-[2-(*tert*-butoxy-carbonylamino)propyl]-1,2,3,4-tetrahydrocyclopent[b]indole (250 mg, 0.59 mmol) in methanol (10 mL) under an atmosphere of Ar at ambient temperature was added hydrogen chloride (4 M in dioxane; 1.4 mL, 5.6 mmol) and then the mixture was stirred for 16 h. Ether (20 mL) was added, and the resultant suspension was cooled (ice-water bath), filtered, and the solid washed
20 with ice-cold ether to afford the product (183 mg, 89%) as a pale turquoise powder. Data for (S)-1-(8-benzyloxy-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine, hydrochloride are included below in Table 6.

Method D(ii): Deprotection using Potassium tert-Butoxide

25

Example 36: (R)-1-(7-Fluoro-1,2,3,4-tetrahydro-6-methoxy-cyclopent[b]indol-4-yl)-2-propylamine, hemifumarate

- To a stirred solution of (R) 4-[2-(*tert*-butoxy-carbonylamino)propyl]-7-fluoro-1,2,3,4-
30 tetrahydro-6-methoxy-cyclopent[b]indole (0.405 g, 1.12 mmol) in methyl sulfoxide (10 mL), under argon at 0 °C was added potassium *tert*-butoxide (0.126 g, 1.12 mmol) portionwise over 4 min. The reaction was stirred under argon at room temperature for 20 h, poured into ice/water (2:1, 150 mL) and stirred until all the ice had melted. The aqueous

suspension was extracted with ethyl acetate (2 x 50 mL). The combined organic extracts were successively washed with water (2 x 20 mL), brine (20 mL) then dried (MgSO₄), filtered and concentrated *in vacuo*. The residue was dissolved in hot 2-propanol (5 mL) and added dropwise to a stirred solution of fumaric acid (0.12 g, 1 mmol) in hot 2-propanol (5 mL). The mixture was cooled to 0 °C, diluted with ether (50 mL) and filtered. The filter-cake was washed (ice-cold 2-propanol, ether) and dried *in vacuo* to yield the hemifumarate as an off-white solid (0.27 g, 75%). Data for (*R*)-1-(7-fluoro-1,2,3,4-tetrahydro-6-methoxy-cyclopent[*b*]indol-4-yl)-2-propylamine, hemifumarate are included in Table 6 below.

10

Method D(iii): Deprotection using Trifluoroacetic Acid


Example 45: (*S*)-1-(3,4-Dihydro-1-oxo-2H-hydrocyclopent[*b*]indol-4-yl)-2-propylamine hydrochloride

15

A stirred solution of (*S*) 4-[2-(*tert*-butoxy-carbonylamino)propyl]-3,4-dihydro-1-oxo-2H-cyclopent[*b*]indole (0.1 g, 0.3 mmol) in dichloromethane (5 mL) was cooled to 0 °C (ice). Trifluoroacetic acid (2 mL, 26 mmol) was added dropwise to the mixture and stirring was continued at 0 °C for 5 h. The mixture was poured onto ice-water (10 mL), basified (pH 8-9) using aqueous sodium hydroxide solution (2 N) then extracted with dichloromethane (2 x 10 mL). The organic extracts were combined, dried (MgSO₄), evaporated to dryness then dissolved in methanol – dichloromethane (1:9, 10 mL), treated with ethereal hydrogen chloride solution (1 M, 1 mmol) and concentrated *in vacuo* to give the title compound as a white solid (0.057 g, 72%). Data for Example 45 are listed below in Table 6.


25

Table 6: Indole-propylamines of formula (I) synthesised using General Method D




In this structural formula, there may be an additional double bond in the 5- or 6-membered ring fused to the indole ring. In Table 6 below, the substituents R_4 to R_7 are hydrogen unless otherwise stated (see column 2). In Table 6 below, the stereochemistry at the side chain is indicated in column 3.

5

Example	Substitution pattern	n	Yield (method)	Data
2	R ₆ =F	1 (S)	63% (i)	Fumarate. m.p. 161-162 °C; Found: C, 61.33; H, 6.11; N, 8.05%. C ₁₈ H ₂₁ FN ₂ O ₄ ·0.25H ₂ O requires: C, 61.27; H, 6.14; N, 7.94%.
3	R ₅ =Cl	2 (S)	84% (i)	Fumarate. m.p. 205 °C (dec.); Found: C, 59.50; H, 6.13; N, 7.23%. C ₁₉ H ₂₃ ClN ₂ O ₄ ·0.25H ₂ O requires: C, 59.53; H, 6.18; N, 7.31%.
4	R ₇ =Cl	1 (S)	25% (i)	Fumarate. m.p. 172-173 °C (dec.); Found: C, 58.63; H, 5.69; N, 7.44%. C ₁₄ H ₁₇ ClN ₂ ·1.1C ₄ H ₄ O ₄ requires: C, 58.71; H, 5.73; N, 7.44%.
5	R ₅ =Cl	1 (S)	17% (ii)	Fumarate. m.p. 175-180 °C (dec.); Found: C, 59.01; H, 5.91; N, 7.34%. C ₁₈ H ₂₁ ClN ₂ O ₄ requires: C, 59.26; H, 5.80; N, 7.67%.
6	R ₅ = Cl; n in the above formula is not applicable; the compound contains an S-heteroatom:  Stereochemistry is (S)	2% (ii)	Hemifumarate. m.p. 189-192 °C; NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.64 (1H, d, <i>J</i> 2 Hz), 7.45 (1H, d, <i>J</i> 8.5 Hz), 7.04 (1H, dd, <i>J</i> 2, 8.5 Hz), 6.47 (1H, s), 4.11 (1H, q, <i>J</i> 7 Hz), 4.04 (1H, q, <i>J</i> 7 Hz), 3.81 (2H, s), 3.36 (2H, m), 3.08 – 2.91 (4H, m), 1.04 (3H, d, <i>J</i> 6.5 Hz).	

7	R ₅ =Br	1 (S)	35% (i)	Hemifumarate. NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.71 (1H, d, <i>J</i> 2 Hz), 7.30 (1H, d, <i>J</i> 8.5 Hz), 7.11 (1H, dd, <i>J</i> 8.5, 2 Hz), 6.46 (1H, s), 4.08 (1H, dd, <i>J</i> 14.5, 6.5 Hz), 3.98 (dd, <i>J</i> 14.5, 7 Hz), 3.35 (4H, m), 2.86 (2H, m), 2.76 (2H, m), 2.48 (2H, quint., <i>J</i> 7 Hz), 1.05 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (4.38 min).
8	R ₅ =Br	2 (S)	51% (i)	Hemifumarate. NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.73 (1H, d, <i>J</i> 1.5 Hz), 7.34 (1H, d, <i>J</i> 8.5 Hz), 7.12 (1H, dd, <i>J</i> 8.5, 1.5 Hz), 6.48 (1H, s), 4.13 (1H, dd, <i>J</i> 14.5, 6.5 Hz), 4.04 (1H, dd, <i>J</i> 14.5, 7.5 Hz), 3.41 (1H, obs. sextet, <i>J</i> 7 Hz), 2.80-2.68 (2H, m), 2.63 (2H, m), 1.82 (2H, m), 1.79 (2H, m), 1.06 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (5.17 min).
9	R ₆ =Cl	2 (S)	78% (i)	Fumarate. NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.49 (1H, d, <i>J</i> 8.5 Hz), 7.42 (1H, d, <i>J</i> 2 Hz), 7.08 (1H, dd, <i>J</i> 8.5, 2 Hz), 6.50 (2H, s), 4.23 (1H, dd, <i>J</i> 14.5, 6.5 Hz), 4.11 (1H, dd, <i>J</i> 14.5, 8.5 Hz), 3.47 (1H, obs. sextet, <i>J</i> 7 Hz), 2.81-2.66 (2H, m), 2.65-2.59 (1H, m), 1.91-1.83 (2H, m), 1.83-1.74 (2H, m), 1.07 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (5.10 min).


10	R ₆ =Cl	1 (S)	62% (i)	Fumarate. NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.49 (1H, d, <i>J</i> 8.5 Hz), 7.39 (1H, d, <i>J</i> 2 Hz), 7.06 (1H, dd, <i>J</i> 8.5, 2 Hz), 6.50 (2H, s), 4.25 (1H, dd, <i>J</i> 14.5, 6.5 Hz), 4.09 (1H, dd, <i>J</i> 14.5, 8 Hz), 3.47 (1H, obs. sextet, <i>J</i> 7 Hz), 2.92 (1H, dd, <i>J</i> 15.5, 7 Hz), 2.84 (1H, dd, <i>J</i> 15.5, 7.5 Hz), 2.76 (2H, t, <i>J</i> 7 Hz), 2.47 (2H, quint, <i>J</i> 7 Hz), 1.09 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (4.48 min).
11	R ₅ =OMe	1 (S)	73% (i)	Fumarate. m.p. 182 °C; Found: C, 63.37; H, 6.75; N, 7.76%. C ₁₉ H ₂₄ N ₂ O ₅ requires: C, 63.32; H, 6.71; N, 7.77%.
12	R ₇ =OMe	1 (S)	75% (i)	Fumarate. NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.05 (1H, d, <i>J</i> 8 Hz), 6.96 (1H, t, <i>J</i> 8 Hz), 6.51 (1H, d, <i>J</i> 8 Hz), 6.50 (2H, s), 4.24 (1H, dd, <i>J</i> 14.5, 6 Hz), 4.04 (1H, 14.5, 8 Hz), 3.82 (3H, s), 3.47 (1H, sextet, <i>J</i> 7 Hz), 2.90-2.75 (4H, m), 2.45 (2H, quint., <i>J</i> 7 Hz), 1.08 (3H, d, <i>J</i> 6.5 Hz), HPLC: [Supelcosil ABZ+ 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 96% (2.90 min).
13	R ₄ =R ₅ =Cl	1 (S)	84% (i)	Hydrochloride. m.p. 288-291 °C; Found: C, 52.84; H, 5.38; N, 8.76; Cl, 33.48%. C ₁₄ H ₁₇ Cl ₃ N ₂ requires: C, 52.60; H, 5.36; N, 8.76; Cl, 33.27%.
14		1 (S)	90% (i)	Hydrochloride. m.p. 233 °C (ethyl acetate); Found: 65.29; H, 7.52; N, 10.81%. C ₁₄ H ₁₈ N ₂ .Hydrochloride.0.375H ₂ O requires: C, 65.30; H, 7.73; N, 10.88%.

15	$R_5 = R_7 = \text{Cl}$; n in the above formula is not applicable; the compound contains an S-heteroatom  Stereochemistry is (S)	74% (i)	Hydrochloride. m.p. 316-322 °C (ethyl acetate); Found: C, 44.54; H, 3.78; N, 7.84%. $\text{C}_{13}\text{H}_{12}\text{Cl}_2\text{N}_2\text{S} \cdot \text{Hydrochloride} \cdot \text{H}_2\text{O}$ requires: C, 44.15; H, 4.27; N, 7.92%.
16	$R_5 = \text{Cl}$ $R_6 = \text{F}$	1 (S)	15% (iii) NMR (400MHz, DMSO- d_6) δ_{H} 1.18 (3H, d, J 6.53Hz), 2.46 (2H, m), 2.73 (2H, m), 2.78-2.95 (2H, m), 3.57 (1H, m), 4.15 (1H, dd, J 7.53Hz, 14.56Hz), 4.36 (1H, dd, J 6.53Hz, 14.05Hz), 7.33 (1H, d, J 9.54Hz), 7.80 (1H, d, J 6.53Hz), 8.27 (3H, br s); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 96% (4.28 min).
17	$R_5 = \text{CF}_3$	1 (S)	94% (i) Hydrochloride. m.p. 270-274 °C (ethyl acetate); Found: C, 56.31; H, 5.83; N, 8.66%. $\text{C}_{15}\text{H}_{17}\text{F}_3\text{N}_2 \cdot \text{HCl}$ requires: C, 56.52; H, 5.69; N, 8.78%.
18	$R_7 = \text{Cl}$ $R_6 = \text{F}$	1 (S)	25% (i) Hydrochloride. m.p. 252-253 °C (ether); Found: C, 54.12; H, 5.60; N, 8.91%. $\text{C}_{14}\text{H}_{17}\text{Cl}_2\text{FN}_2 \cdot 0.5\text{H}_2\text{O}$ requires: C, 53.86; H, 5.81; N, 8.97%.
19	$R_5 = R_6 = \text{Cl}$	1 (S)	93% (i) Hydrochloride. m.p. 292-295 °C (ethyl acetate); Found: C, 52.20; H, 5.29; N, 8.63%. $\text{C}_{14}\text{H}_{16}\text{Cl}_2\text{N}_2 \cdot \text{HCl} \cdot 0.25\text{H}_2\text{O}$ requires: C, 51.87; N, 5.44; N, 8.64%.

20	$R_6=OMe$	1 (S)	79% (i)	Hydrochloride. m.p. 260 °C (dec.); NMR (400 MHz, DMSO- d_6) δ_H 8.37 (3H, m, NH_3), 7.40 (1H, d, J 8.5 Hz), 6.88 (1H, d, J 2.5 Hz), 6.71 (1H, dd, J 8.5, 2.5 Hz), 4.36 (1H, dd, J 14.5, 6 Hz), 4.11 (1H, dd, J 14.5, 8 Hz), 3.76 (3H, s), 3.54 (1H, m), 3.39 (1H, m), 2.94-2.78 (2H, m), 2.75 (2H, t, J 7 Hz), 2.47 (2H, quintet, J 7 Hz), 1.16 (3H, d, J 6.5 Hz).
21	$R_7=CF_3$	1 (S)	59% (i)	Hydrochloride. m.p. 238-242 °C; NMR (400 MHz, DMSO- d_6) δ_H 8.40 (3H, m, NH_3), 7.89 (1H, d, J 8 Hz), 7.39 (1H, d, J 8 Hz), 7.23 (1H, t, J 7 Hz), 4.47 (1H, dd, J 14.5, 6.5 Hz), 4.28 (1H, dd, J 14.5, 7.5 Hz), 3.61 (1H, m), 3.39 (1H, m), 3.04-2.86 (2H, m), 2.82 (2H, t, J 7 Hz), 2.50 (2H, quint., J 7 Hz), 1.22 (3H, d, J 6.5 Hz).
22	$R_6=R_7=Cl$	1 (S)	74% (i)	Hydrochloride. m.p. 243-248 °C (ethyl acetate); Found: C, 51.20; H, 5.30; N, 8.28%. $C_{14}H_{16}Cl_2N_2 \cdot HCl \cdot 0.5H_2O$ requires: C, 51.16; H, 5.52; N, 8.52%.
23	$R_7=OBn$	1 (S)	86%	Hydrochloride. NMR (400 MHz, DMSO- d_6) δ_H 8.35 (3H, m, NH_3), 7.50 (2H, d, J 7.5 Hz), 7.42 (2H, t, J 7.5 Hz), 7.33 (1H, t, J 7.5 Hz), 7.12 (1H, d, J 8.5 Hz), 6.98 (1H, t, J 8 Hz), 6.63 (1H, d, J 8 Hz), 5.19 (2H, s), 4.36 (1H, dd, J 14.5, 6 Hz), 4.13 (1H, dd, J 14.5, 8 Hz), 3.56 (1H, m), 3.44 (1H, m), 2.96-2.77 (4H, m), 2.48 (2H, quint., J 7 Hz), 1.17 (3H, d, J 6.5 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (5.15 min).

24	$R_7=O^iPr$	1 (S)	43% (iii)	Fumarate. m.p. 189 °C (dec.); NMR (400 MHz, DMSO- d_6) δ_H 7.02 (1H, d, J 8.0 Hz), 6.93 (1H, t, J 7.4 Hz), 6.49 (2H, s), 4.58 (1H, quint, J 6.0 Hz), 4.17 (1H, dd, J 14.4, 6.2 Hz), 4.00 (1H, dd, J 14.4, 7.8 Hz), 3.44 (1H, obs sextet, J 6.7 Hz), 2.91-2.76 (4H, m), 2.44 (2H, obs quint, J 7.0 Hz), 1.30 (6H, d, J 6.0 Hz), 1.08 (3H, d, J 6.5 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 98.4% (3.33 min); Found C, 65.31; H, 7.36; N, 7.36%. $C_{21}H_{28}N_2O_5$ requires: C, 64.93; H, 7.26; N, 7.21%
25	$R_5=O^iPr$	1 (S)	35% (ii)	Hemifumarate. m.p. 163 °C (dec.); NMR (400 MHz, DMSO- d_6) δ_H 7.19 (1H, d, J 8.6 Hz), 7.02 (1H, d, J 2.0 Hz), 6.61 (1H, dd, J 8.6, 2.0 Hz), 6.46 (1H, s), 4.61 (1H, quint, J 6.0 Hz), 4.07 (1H, dd, J 14.3, 6.2 Hz), 3.92 (1H, dd, J 14.3, 7.5 Hz), 3.34 (1H, q, J 6.7 Hz), 2.88-2.78 (2H, m), 2.73 (2H, obs t, J 6.8 Hz), 2.45 (2H, obs quint, J 6.9 Hz), 1.27 (6H, d, J 6.0 Hz), 1.04 (3H, d, J 6.7 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 97% (3.35 min); Found C, 69.04; H, 7.92; N, 8.42%. $C_{19}H_{26}N_2O_3$ requires: C, 69.07; H, 7.93; N, 8.47%.
26	$R_5=R_7=Cl$	1 (S)	99% (i)	Hydrochloride. m.p. 258 °C (ethyl acetate); Found: C, 52.02; H, 5.28; N, 8.53%. $C_{14}H_{16}Cl_2N_2 \cdot HCl \cdot 0.25H_2O$ requires: C, 51.87; H, 5.44; N, 8.64%.

27	R ₅ =EtS	1 (S)	46%	Hydrochloride. m.p. 115-119 °C; NMR (400MHz, DMSO- <i>d</i> ₆) δ _H 1.17 (6H, m), 2.46 (2H, m), 2.73 (2H, m), 2.85, (2H, m), 2.95 (2H, q, <i>J</i> 7.53Hz), 3.53 (1H, m), 4.14 (1H, dd, <i>J</i> 7.53, 14.05), 4.36 (1H, dd, <i>J</i> 6.53Hz, 14.05Hz), 7.02 (1H, d, <i>J</i> 8.53Hz), 7.29 (1H, d, <i>J</i> 8.53Hz), 7.56 (1H, br s), 8.34 (3H, br s).
28	R ₄ =OCF ₃	1 (S)	70% (i)	Hydrochloride. m.p. 281 °C (dec); NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 8.45 (3H, m, NH ₃), 7.38 (1H, dd, <i>J</i> 7.2, 1.5 Hz), 7.11 –7.05 (2H, m), 4.37 (1H, dd, <i>J</i> 14.8, 7.1 Hz), 4.29 (1H, dd, <i>J</i> 14.8, 6.9 Hz), 3.52 (1H, q, <i>J</i> 6.5 Hz), 3.00 (1H, obs quint, <i>J</i> 7.5 Hz), 2.89 (1H, obs quint, <i>J</i> 6.9 Hz), 2.81-2.78 (2H, m), 2.53-2.47 (2H, m), 1.16 (3H, d, <i>J</i> 6.7 Hz); HPLC: [Supelcosil ABZ+; 1.0 ml/min, methanol-10mM aqueous ammonium acetate solution (80:20)] 99.8% (3.80 min).
29	R ₆ =OCF ₃	1	95% (i)	Hydrochloride. m.p. 166-169 °C; NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 8.43 (3H, m, NH ₃), 7.66 (1H, d, <i>J</i> 9 Hz), 7.33 (1H, d, <i>J</i> 1.5 Hz), 7.05 (1H, dd, <i>J</i> 9, 1.5 Hz), 4.44 (1H, dd, <i>J</i> 14.5, 6.5 Hz), 4.21 (1H, dd, <i>J</i> 14.5, 8 Hz), 3.59 (1H, m), 3.00-2.81 (2H, m), 2.78 (2H, t, <i>J</i> 7 Hz), 2.49 (2H, quint, <i>J</i> 7 Hz), 1.20 (3H, d, <i>J</i> 6.5 Hz).
30		1 (R)	92% (i)	Hydrochloride. m.p. 225-231 °C (dec.); Found: C, 65.37; H, 7.51; N, 10.78%. C ₁₄ H ₁₈ N ₂ .HCl.0.33H ₂ O requires: C, 65.49; H, 7.33; N, 10.91%.
31	R ₅ =F	1 (S)	40% (i)	Hydrochloride. m.p. 215 °C (ether); Found: C, 60.38; H, 6.58; N, 9.85%. C ₁₄ H ₁₇ FN ₂ .HCl.0.5H ₂ O requires: C, 60.54; H, 6.53; N, 10.09%.


32	n in the above formula is not applicable; the compound contains an S-heteroatom:  Stereochemistry is (S)	97% (i)	Hydrochloride. m.p. 289-293 °C (ethyl acetate); Found: C, 58.57; H, 5.77; N, 10.49%. C ₁₃ H ₁₄ N ₂ S.HCl requires: C, 58.53; H, 5.67; N, 10.50%.	
33	R ₅ =R ₆ =F	1	21% (i)	(Free-base purified by column chromatography, [SiO ₂ ; ethyl acetate – methanol – ammonium hydroxide (92:7:1)]. Hydrochloride. m.p. 249-250 °C; NMR (400MHz, DMSO- <i>d</i> ₆) δ _H 1.18 (3H, d, <i>J</i> 6.53Hz), 2.45 (2H, m), 2.72 (2H, m), 2.77-2.94 (2H, m), 3.55 (1H, br s), 4.13 (1H, dd, <i>J</i> 8.03Hz, 15.06Hz), 4.35 (1H, dd, <i>J</i> 6.53Hz, 14.56Hz), 7.33 (1H, dd, 8.03Hz, 11.04Hz), 7.72 (1H, dd, <i>J</i> 7.03Hz, 12.05Hz), 8.35 (3H, br s).
34	R ₇ =Cl R ₆ =Me	1 (S)	62% (i)	Hydrochloride. m.p. 250+ °C (dec.); NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 8.14 (3H, m, -NH ₃), 7.36 (1H, d, <i>J</i> 8.5 Hz), 7.02 (1H, d, <i>J</i> 8.5 Hz), 4.30 (1H, dd, <i>J</i> 6.5, 15 Hz), 4.14 (1H, dd, <i>J</i> 7.5, 14.5 Hz), 3.57 (1H, m), 2.98 (2H, app. t, <i>J</i> 7 Hz), 2.92-2.80 (2H, m), 2.48 (2H, quint., <i>J</i> 7 Hz), 2.38 (3H, s), 1.17 (3H, d, <i>J</i> 6.5 Hz).
35	R ₇ =Cl R ₆ =Me	1 (R)	44% (i)	Hydrochloride. m.p. 250+ °C (dec.); NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 8.28 (3H, m, -NH ₃), 7.38 (1H, d, <i>J</i> 8.5 Hz), 7.01 (1H, d, <i>J</i> 8.5 Hz), 4.34 (1H, dd, <i>J</i> 6.5, 14.5 Hz), 4.15 (1H, dd, <i>J</i> 1.5, 14.5 Hz), 3.56 (1H, m), 2.98 (1H, app. t, <i>J</i> 7 Hz), 2.93-2.80 (2H, m), 2.48 (2H, quint., <i>J</i> 7 Hz), 2.38 (3H, s), 1.17 (3H, d, <i>J</i> 6.5 Hz).

36	R ₆ =F R ₅ =OMe	1 (R)	75% (ii)	Fumarate. NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.30 (1H, d, <i>J</i> 7.4 Hz), 7.14 (1H, d, <i>J</i> 12.1 Hz), 6.51 (2H, s), 4.26 (1H, dd, <i>J</i> 14.6, 5.8 Hz), 4.07 (1H, dd, <i>J</i> 14.6, 7.0 Hz), 3.88 (3H, s, MeO), 3.52 (1H, br s), 2.91-2.78 (2H, m), 2.74-2.68 (2H, m), 2.45 (2H, obs quint, <i>J</i> 7.1 Hz), 1.11 (3H, d, <i>J</i> 6.3 Hz); HPLC: [Xterra; 2.0 ml/min, methanol-10mM aqueous ammonium acetate solution, gradient elution (50:50) to (80:20) over the first 4 min, then (80:20)] 97% (2.40 min).
37	R ₆ =F R ₅ =OMe	1 (S)	100% (ii)	Fumarate. m.p. 213 °C (dec.); Found: C, 59.99; H, 6.24; N, 7.08%. C ₁₉ H ₂₃ N ₂ O ₅ F requires: C, 60.31; H, 6.13; N, 7.40%.
39	R ₅ =Cl R ₆ =F	1 (R)	58% (i)	Fumarate. NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.75 (1H, d, <i>J</i> 6.5 Hz), 7.32 (1H, d, <i>J</i> 10 Hz), 6.51 (2H, s), 4.20 (1H, dd, <i>J</i> 6.5, 14.5 Hz), 4.07 (1H, dd, <i>J</i> 1, 14.5 Hz), 3.46 (1H, m), 2.96-2.80 (2H, m), 2.75 (2H, app. t, <i>J</i> 7 Hz), 2.47 (2H, quint., <i>J</i> 7 Hz), 1.11 (3H, d, <i>J</i> 6.5 Hz); HPLC: [Xterra; 2.0 ml/min, methanol-10mM aqueous ammonium acetate solution, gradient elution (50:50) to (80:20) over the first 4 min, then (80:20)] 96% (4.72 min).
40	R ₇ =Cl R ₆ =F	1 (R)	98% (i)	Hydrochloride. m.p. 261-264 °C (ethyl acetate); Found: C, 53.92; H, 5.61; N, 8.97%. C ₁₄ H ₁₆ ClFN ₂ .HCl.0.5H ₂ O requires: C, 53.86; H, 5.49; N, 8.97%.
41	R ₇ =Br	1 (S)	97% (i)	Hydrochloride. m.p. 246-252 °C (ethyl acetate); Found: C, 49.55; H, 5.45; N, 8.17%. C ₁₄ H ₁₇ BrN ₂ .HCl.0.5H ₂ O requires: C, 49.65; H, 5.36; N, 8.27%.

42	R ₆ =F R ₇ =OMe	1 (S)	37% (ii)	Hemifumarate. NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 7.14 (1H, dd, <i>J</i> 8.8, 3.4 Hz), 6.90 (1H, dd, <i>J</i> 11.7, 8.8 Hz), 6.47 (1H, s), 4.07 (1H, dd, <i>J</i> 14.5, 5.8 Hz), 3.99-3.94 (1H, m), 3.90 (3H, s, MeO), 3.35 (1H, br s), 2.90-2.79 (4H, m), 2.47 (2H, obs quint, <i>J</i> 7.1 Hz), 1.04 (3H, d, <i>J</i> 5.4 Hz); HPLC: [Xterra; 2.0 ml/min, methanol-10mM aqueous ammonium acetate solution, gradient elution (50:50) to (80:20) over the first 4 min, then 80:20] 99.4% (1.61 min).
43	R ₄ =Cl	1 (R)	37% (i)	m.p. 299-302 °C (2-propanol); Found: C, 58.79; H, 6.52; N, 9.48, Cl, 24.51%. C ₁₄ H ₁₈ N ₂ Cl ₂ requires: C, 58.96; H, 6.36; N, 9.82; Cl, 24.86%.
44	R ₄ =Cl	1 (S)	33% (i)	m.p. 296-299 °C (2-propanol); NMR (400 MHz, DMSO- <i>d</i> ₆) δ _H 8.45 (3H, m, NH ₃), 7.32 (1H, dd, <i>J</i> 1, 8 Hz), 7.07 (1H, dd, <i>J</i> 1, 8 Hz), 6.99 (1H, t, <i>J</i> 8 Hz), 4.62 (1H, dd, <i>J</i> 6.5, 14.5 Hz), 4.47 (1H, dd, <i>J</i> 6.5, 14.5 Hz), 3.62 (1H, m), 3.32 (1H, m), 3.00-2.80 (2H, m), 2.76 (2H, m), 2.50-2.43 (2H, m), 1.13 (3H, d, <i>J</i> 7 Hz).
45	1-ketone	1 (S)	72% (iii)	NMR (400MHz, DMSO- <i>d</i> ₆) δ _H 1.29 (3H, d, <i>J</i> 6.53 Hz), 2.85 (2H, m), 3.05-3.13 (1H, m), 3.18-3.27 (1H, m), 3.71 (1H, m), 4.38 (1H, dd, <i>J</i> 6.53 Hz, 14.56Hz), 4.55 (1H, dd, <i>J</i> 7.03 Hz, 15.06 Hz), 7.25 (1H, m), 7.32 (1H, dt, <i>J</i> 1.0Hz, 7.03 Hz), 7.72, 1H, d, <i>J</i> 7.53 Hz), 7.77 (1H, d, <i>J</i> 8.03 Hz), 8.58 (3H, br s); HPLC: [Xterra; 2.0 ml/min, methanol-10mM aqueous ammonium acetate solution (50:50)] 98% (2.11 min).

CLAIMS

1. A chemical compound of formula (I):



(I)

wherein:

R₁ and R₂ are independently selected from hydrogen and alkyl;

R₃ is alkyl;

R₄, R₆ and R₇ are independently selected from hydrogen, halogen, hydroxy, alkyl, aryl, amino, alkylamino, dialkylamino, alkoxy, aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl, nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl;

R₅ is selected from hydrogen, halogen, hydroxy, alkyl, aryl, amino, alkylamino, dialkylamino, alkoxy, aryloxy, alkylthio, alkylsulfoxyl, alkylsulfonyl, nitro, carbonitrile, carbo-alkoxy, carbo-aryloxy and carboxyl; and

A is a 5- or 6-membered partially unsaturated or aromatic heterocyclic ring or a 5- or 6- membered partially unsaturated carbocyclic ring,

wherein if A is a 6-membered partially unsaturated carbocyclic ring then at least one of R₄ to R₇ is other than hydrogen,

and pharmaceutically acceptable salts, addition compounds and prodrugs thereof.

2. A compound according to claim 1 wherein R₁ and R₂ are selected from hydrogen and lower alkyl.

3. A compound according to claim 1 wherein R₁ and R₂ are hydrogen.

4. A compound according to claim 1, 2 or 3 wherein R₃ is lower alkyl.

5. A compound according to claim 1, 2 or 3 wherein R₃ is methyl.
6. A compound according to any preceding claim wherein R₄ is selected from hydrogen, halogen, alkyl and alkoxy.
- 5 7. A compound according to any preceding claim wherein R₄ is hydrogen.
8. A compound according to any preceding claim wherein R₆ is selected from hydrogen and halogen.
- 10 9. A compound according to any preceding claim wherein R₇ is selected from hydrogen, halogen and alkoxy.
10. A compound according to any preceding claim wherein A is a 5- membered ring.
- 15 11. A compound according to any preceding claim wherein A is partially unsaturated.
12. A compound according to any preceding claim wherein A contains a heteroatom selected from N, O and S.
- 20 13. A compound according to any of claims 1 to 9 wherein A is a 5- membered partially unsaturated carbocyclic ring, a 5- membered partially unsaturated or aromatic heterocyclic ring or a 6- membered partially unsaturated carbocyclic ring.
- 25 14. A compound according to any of claims 1 to 9 wherein A is selected from cyclopentenyl, cyclohexenyl, thiacyclohexenyl and thienyl.
15. A compound according to claim 1 which is selected from (S)-1-(7,8-difluoro-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine, (S)-1-(7-fluoro-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine, (S)-1-(8-chloro-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine, (S)-1-(6-methoxy-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine, (S)-1-(7-fluoro-6-methoxy-1,2,3,4-tetrahydrocyclopent[b]indol-4-yl)-2-propylamine, (S)-1-(7-fluoro-8-
- 30

methoxy-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine, (*S*)-1-(8-chloro-7-fluoro-1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine, (*S*)-1-(1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine, (*R*)-1-(1,2,3,4-tetrahydrocyclopent[*b*]indol-4-yl)-2-propylamine.

- 5
16. A compound of formula (I) as set out in any one of claims 1 to 15 for use in therapy.
- 10 17. The use of a compound of formula (I) as set out in any of claims 1 to 15 in the manufacture of a medicament for the treatment of disorders of the central nervous system; damage to the central nervous system; cardiovascular disorders; gastrointestinal disorders; diabetes insipidus, and sleep apnea.
- 15 18. A use according to claim 17 wherein the disorders of the central nervous system are selected from depression, atypical depression, bipolar disorders, anxiety disorders, obsessive-compulsive disorders, social phobias or panic states, sleep disorders, sexual dysfunction, psychoses, schizophrenia, migraine and other conditions associated with cephalic pain or other pain, raised intracranial pressure, epilepsy, personality disorders, age-related behavioural disorders, behavioural disorders
- 20 associated with dementia, organic mental disorders, mental disorders in childhood, aggressivity, age-related memory disorders, chronic fatigue syndrome, drug and alcohol addiction, obesity, bulimia, anorexia nervosa and premenstrual tension.
- 25 19. A use according to claim 17 wherein the damage to the central nervous system is by trauma, stroke, neurodegenerative diseases or toxic or infective CNS diseases.
20. A use according to claim 19 wherein said toxic or infective CNS disease is encephalitis or meningitis.
- 30 21. A use according to claim 17 wherein the cardiovascular disorder is thrombosis.
22. A use according to claim 17 wherein the gastrointestinal disorder is dysfunction of gastrointestinal motility.

23. A use according to claim 17 wherein said medicament is for the treatment of obesity.
- 5 24. A use according to any one of claims 17 to 23 wherein said treatment is prophylactic treatment.
25. A method of treatment of any of the disorders set out in claims 17 to 22 comprising administering to a patient in need of such treatment an effective dose of a
10 compound of formula (I) as set out in any one of claims 1 to 15.
26. A method of treatment according to claim 25 wherein said disorder is obesity.
27. A method according to claim 25 or 26 wherein said treatment is prophylactic
15 treatment.
28. A method of preparing a compound of formula (I) as set out in any one of claims 1 to 15.
- 20 29. A pharmaceutical composition comprising a compound of formula (I) as set out in any one of claims 1 to 15 in combination with a pharmaceutically acceptable carrier or excipient.
30. A method of making a composition according to claim 29 comprising combining a
25 compound of formula (I) as set out in any one of claims 1 to 15 with a pharmaceutically acceptable carrier or excipient.

INTERNATIONAL SEARCH REPORT

Inter- national Application No

PCT/GB 00/03011

CLASSIFICATION OF SUBJECT MATTER

7 C07D209/80 C07D495/04 A61K31/403 A61K31/407 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 655 440 A (HOFFMANN LA ROCHE) 31 May 1995 (1995-05-31) cited in the application claims 1-14,16-22 ---	1,4,5, 16-18, 25,28,29
Y	EP 0 657 426 A (HOFFMANN LA ROCHE) 14 June 1995 (1995-06-14) cited in the application claims ---	1,4,5, 16-18, 25,28,29
Y	US 3 329 571 A (L. M. RICE ET AL.) 4 July 1967 (1967-07-04) cited in the application column 1, line 13 - line 19; example 4 --- -/--	1,16-18, 28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

31 October 2000

Date of mailing of the international search report

10/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/03011

(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 930 988 C (FARBENFABRIKEN BAYER AG) 28 July 1955 (1955-07-28) cited in the application examples 1,2,4,6,8-11 ---	1,28
A	US 3 142 678 A (L. M. RICE ET AL.) 28 July 1964 (1964-07-28) cited in the application column 1, line 23 - line 29; claims 1,3,8 ---	1,16,17, 28
A	US 2 541 211 A (J. W. CUSIC ET AL.) 13 February 1951 (1951-02-13) cited in the application examples 5,6 ---	1,28
A	US 2 687 414 A (J. W. CUSIC) 24 August 1954 (1954-08-24) cited in the application column 3, line 69 - line 72; examples 19,21 ---	1,16,28
A	EP 0 700 905 A (HOFFMANN LA ROCHE) 13 March 1996 (1996-03-13) cited in the application claims ---	1,16-18, 25,28,29
A	WO 98 30548 A (TSUKAMOTO SHIN ICHI ;KUBOTA HIDEKI (JP); MAENO KYOICHI (JP); SHIMA) 16 July 1998 (1998-07-16) cited in the application examples 27-32,46,35,36,47,49,50,51 -----	1,16,17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/03011

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0655440 A	31-05-1995	AU 685841 B	29-01-1998
		AU 7583794 A	11-05-1995
		BR 9404203 A	04-07-1995
		CA 2132883 A	23-04-1995
		CN 1105988 A	02-08-1995
		CZ 9402604 A	18-10-1995
		FI 944969 A	23-04-1995
		HU 70848 A	28-11-1995
		IL 111314 A	17-08-1999
		JP 2638752 B	06-08-1997
		JP 7149723 A	13-06-1995
		NO 943999 A	24-04-1995
		NZ 264713 A	28-05-1996
		PL 305543 A	02-05-1995
		RU 2136662 C	10-09-1999
		US 5494928 A	27-02-1996
		ZA 9408094 A	24-04-1995
EP 0657426 A	14-06-1995	AT 168675 T	15-08-1998
		AU 680543 B	31-07-1997
		AU 7583894 A	11-05-1995
		BR 9404205 A	04-07-1995
		CA 2132887 A	23-04-1995
		CN 1105989 A,B	02-08-1995
		CZ 9402603 A	12-07-1995
		DE 59406489 D	27-08-1998
		DK 657426 T	26-04-1999
		ES 2120551 T	01-11-1998
		FI 944970 A	23-04-1995
		HU 70443 A,B	30-10-1995
		JP 2638751 B	06-08-1997
		JP 7149725 A	13-06-1995
		NO 944000 A	24-04-1995
		NZ 264711 A	26-07-1996
		PL 305544 A	02-05-1995
		RU 2128649 C	10-04-1999
		US 5646173 A	08-07-1997
		ZA 9408093 A	05-06-1995
US 3329571 A	04-07-1967	CH 470384 A	31-03-1969
		DE 1445416 A	23-01-1969
		DK 120748 B	12-07-1971
		DK 120541 B	14-06-1971
		DK 116282 B	29-12-1969
		NL 133522 C	
		NL 284655 A	
DE 930988 C		NONE	
US 3142678 A	28-07-1964	NONE	
US 2541211 A	13-02-1951	NONE	
US 2687414 A	24-08-1954	NONE	
EP 0700905 A	13-03-1996	AU 691310 B	14-05-1998
		AU 2838295 A	22-02-1996
		BR 9503630 A	28-05-1996

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/03011

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0700905 A		CA 2153937 A	13-02-1996
		CN 1131666 A,B	25-09-1996
		CZ 9502013 A	13-03-1996
		FI 953827 A	13-02-1996
		HU 72066 A	28-03-1996
		JP 2755560 B	20-05-1998
		JP 8059623 A	05-03-1996
		NO 953162 A	13-02-1996
		NZ 272731 A	25-09-1996
		PL 309974 A	19-02-1996
		TR 960124 A	21-06-1996
		US 5561150 A	01-10-1996
		ZA 9506553 A	12-02-1996
WO 9830548 A	16-07-1998	AU 5343298 A	03-08-1998

THIS PAGE BLANK (USPTO)

THIS PAGE BLANK (USPTO)

THIS PAGE BLANK (USPTO)